CT Angio in Coarctation of Aorta

Gojwari T, Feroze S, Irfan Robbani, Singh M, Farooq Mir M, Parvez Nazir, Nisar A Wani, Shafeeq A, Naseer Khan and Ajaz Hakeem

Department of Radiodiagnosis, Sher-i-Kashmir Institute of Medical Sciences, Srinagar.

Definition

It is a congenital narrowing in the region of isthmus, more common in males and rare in blacks.

Introduction/Discussion

It is generally of two types. Localized or Adult or Postductal type: It is most common type where there is short segment of narrowing close to the ligamentum arteriosum. Coexisting cardiac anomalies are uncommon.

Tubular hypoplasia or Infantile or Diffuse type: there is hypoplasia of relatively longer segment of the aortic arch after the origin of the brachio-cephalic trunk. Coexisting cardiac anomalies are common. Associated anomalies are Bicuspid aortic valve, Cardiac shunts, Turner's syndrome and Cerebral artery aneurysms.

Collateral circulation distal to the Coarctation is through Subclavian arteries and branches—Internal mammary arteries to intercostal, scapular artery to anterior spinal artery and transverse cervical artery to lateral thoracic artery.

Radiology

- Conventional Radiograph
- Rib notching due to pressure erosion by enlarged and tortuous intercostal arteries. It is bilateral but asymmetrical.
- Left ventricular enlargement.
- Figure of “3” appearance due to dilated left subclavian artery above Coarctation, pinching at site of Coarctation and poststenotic dilatation distally.
- Small and flat aortic knuckle.
- CTA (CT Angio) with multi-slice scanners especially

64 slice or above gives excellent information and accurately depicts the site (Fig. 1), degree and the size of narrowing, collateral formations (Fig. 2 & Fig. 3). Multiplanar imaging and reconstructions give actual visual impression to surgeons.

Figure 1: Saggital reformation showing tight focal narrowing in post ductal region with large vertebral collaterals

Figure 2: Large dilated collaterals around scapula

Figure 3: Dilated Internal mammary arteries

Reference

1. Campbell M. Natural history of Coarctation of aorta. *Br Heart J* 1970;32:633-40.

Reprint request:

Dr. Tariq A. Gojwari

Head, Department of Radiodiagnosis and Imaging
SKIMS, P. Box: 27, Srinagar-190 011