ISSN 0972-110X
eISSN 2582-063X
July-Sept, 2020
Volume 23 Number 3

We are
Crossref
Member

JOURNAL OF MEDICAL SCIENCES

www.jmsskims.org

Indexed with

INDEX COPERNICUS
I N T E R N A T I O N A L

Google
Scholar

Multi disciplinary Peer Reviewed Journal of Medical and Allied Sciences

**An official publication of
Sher-i-Kashmir Institute of Medical Sciences (SKIMS) Srinagar Kashmir India**

Editorial Board

Patron

Prof. A. G. Ahangar

Director & EOSG

Secretary to Govt.

Editor-in-Chief

Shariq R Masoodi (DM)

Prof. Endocrinology, SKIMS Srinagar

Executive Editors

Zafar Amin Shah (Ph.D. Immunology)

Ajaz Ahmad Malik (MS Gen. Surgery)

Ajaz Nabi Koul (MD Internal Medicine)

Associates Editors

Tariq A. Gojwari (MD Radiodiagnosis))

Bashir A. Fomda (MD Microbiology)

Altaf H. Shah (DM Gastroenterology)

Mohammad Iqbal Lone (MD Pathology)

Nayil Khurshid (MCh Neurosurgery)

Rafi Ahmad Jan (MD Internal Medicine)

Adil Hafeez (DNB Plastic Surgery)

Muzafar Maqsood Wani (DM Nephrology)

Bashir Ahmad Dar (MD Anesthesiology)

Assistant Editors

Peerzada Umar Farooq Baba (M.Ch Plastic Surgery)

Syed Mudasir Qadri (MD Internal Medicine)

Suhail Mantoo (MD Internal Medicine)

Mahrukh Hameed Zargar (PhD)

Web Editors

Shabir Ahmed Naikoo (MCA)

Rafia Qadri (MCA)

Editorial Advisory Board

Abdul Q.Haji (U.S.A.)

Altaf Bukhari (KSA)

Anil Sharma (India)

Anil Arora (India)

B. Indira Devi (India)

D.K. Bhasin (India)

Dhananjay Bhat (India)

Dhaval Shukla (India)

Farooq A Wani (India)

Faroque A. Khan (U.S.A.)

Ghulam Jeelani Mufti (U.K)

Gulzar Mufti (U.K)

Imtiyaz Naqash (India)

Javed Latoo (U.K)

M Maqbool Lone (India)

Mehboob Ali Dar (KSA)

Mohammad Azizur Rahman (Bangladesh)

Mushtaq A Wani (U.K)

N A Chowdhary (India)

Nasib Chand Digra (India)

Nupur Pruthi (India)

P. Kar (India)

Pavanasam Velayutham (India)

Raffit Hassan (U.S.A.)

Romesh Kardori (U.S.A.)

S. K. Sharma (India)

S.K. Sarin (India)

Saleem Kamili (U.S.A.)

Shaida Andrabi (U.S.A.)

Sridhar Chitturi (Australia)

Tariq Majid Wani (Qatar)

Vikas.V (India)

Former Editors

A.H. Zargar (1998-2002)

Musthaq A Siddiqi (2003-2007)

Showkat A Zargar (2008-2011)

Parvaiz A Koul (2012-2015)

Aims and Scope

Journal of Medical Sciences, the official journal of Sher-i-Kashmir Institute of Medical Sciences (SKIMS) Srinagar is a peer-reviewed multidisciplinary medical journal which publishes original research articles, reviews, brief communications, case reports and images in Medicine, Surgery, Basic Sciences and their allied specialties. The Journal is published biannually and is available online as well.

Copyright and Photocopying

Papers accepted automatically become the copyright of JMS. Photocopies of single article made for personal use is allowed as per National copyright laws. Permission by the editor/publisher is required for all other photocopying including multiple or systematic copyright, copying for promotional purposes, resales, and all forms of document delivery.

Disclaimer

The papers contained in the journal have been prepared and written by named authors. The statements, opinions, and ideas expressed within these papers are the sole responsibility of these authors and do not imply an opinion on the part of the officers or members of JMS Editorial Board. Therefore, neither SKIMS, nor office bearers, employees or agents of JMS editorial board are responsible for the accuracy or otherwise of any paper, and shall not be liable for any claim, damage or loss arising from the contents of any paper, or any use of which they may be put by any person. Readers are urged to exercise professional caution in undertaking any of the recommendations or suggestions made by the authors.

Printing:

Layout Design: Shah Nawaz A Khosa

Printed at: Zum Zum Printers

Editorial Secretariat

Room No. 11, 3rd Floor
New Faculty Block (Old Library)
Sher-i-Kashmir Institute of Medical Sciences
Post Bag 27, Srinagar 190011, Kashmir (India)
Phone ++91 194-2402029 (Off)
01942401013 Ext. 2064
+919419013289 (Mobile)
Fax: +911942402029, +911942405507
E-mail: editor@jmsskims.org
shariq.masoodi@skims.ac.in

Contents

EDITORIAL

Bacterial Infection in etiology of Pigment Gallstones

1

Rauf Ahmad Wani

REVIEW ARTICLE

Immune checkpoint inhibitor (ICPI) induced Nephrotoxicity An Insight

2

Mohmad Hussian Mir, Tariq Ahmad Bhat, Khalid Parvez Sofi, Imtiyaz Ahmad Wani, Muzafar Maqsood Wani

MINI REVIEW ARTICLE

The Potential Impacts Of Obesity On COVID-19

6

Tasaduk Sultan Khan, Musadiq Sultan Khan

ORIGINAL ARTICLES

Study of Bacterial Aetiogenicity of Pigment Gall Stones by Culture of Nidus of Calculus and correlation between Stone Culture and Bile Culture

8

Mahmood Rashid Shiekh, Tahleel Shera, Shiekh Aamir Rashid, Altaf Hussain Shera, Junaid Ahmad,

Clinical spectrum and outcomes of snakebite patients admitted in a tertiary care hospital.

A prospective observational hospital-based study

13

Sameer Anand, Owais Ahmed Wani, Vijay Kundal

**Incidence and outcome of early post stroke seizures in a tertiary care hospital in New Delhi.....
a prospective observational study**

18

Maqsood Ahmad Dar, Vasundhara Aggarwal Saika, Joy Dev Mukherjee, Mukesh Kumar, Khushbu Goel

Effect of Metformin on TSH Levels

24

Idrees Mubarak, Shariq Rashid Masoodi, Zafar Amin Shah, Syed Mudassar, Saba Moosaraza, Mir Fawad Ahmed

CASE REPORTS

Recurrent Small Gut Fecalomas Causing Obstruction- A Case Report

30

Mudassir Ahmad Khan, Asif Mehraj, Arshad Baba, Mushtaq Laway, Rauf A Wani,

Fazl Q Parry, Nisar A Chowdri

Case of Van-Wyk Grumbach syndrome

33

Vishnu Vasudevan, Arun Viswanath, Riaz Ahmad Misgar

CLINICAL IMAGE

Short RP Tachycardia, Successfully Managed with Radio Frequency Ablation

36

Ajaz Ahmad Lone, Nisar Ahmad Trambo, Syed Bilal

Nasal Reconstruction

38

Peerzada Umar Farooq Baba, Adil Hafeez Wani

LETTERS

Adoption of Telemedicine -need of the hour

39

Sridhar Chitturi

PRACTICE POINT

High Flow Nasal Cannula Oxygenation

41

Tajamul Hussain Shah, Suhail Mantoo, Rafi A Jan

NEWS AND VIEWS

45

Bacterial Infection in etiology of Pigment Gallstones

Gall Stone Disease is widely prevalent in world with Ultrasound studies showing mean prevalence rates of 10-15% in adult European, and of 3-5% of adult African & Asian populations.

Gallstones can be biochemically divided into 3 classes: 1. Pure Cholesterol Gallstones 2. Pigment Gallstones 3. Mixed Gallstones

Majority of gallstones are cholesterol stones. Pigment gallstones are made up of calcium bilirubinate (main Ingredient). They are of brown or black colour depending upon their composition. Brown pigment stones are an important cause of acute cholangitis in Oriental Cholangiohepatitis. Bacterial infection and the deconjugation of bile by bacterial -Glucourindase have been implicated as etiological factors in formation of brown pigment stones but controversy exists regarding their primary importance and causal relationship.

The stones have been analyzed by various methods, besides stone cultures, including by chemical microscopy, Infrared Spectrophotometry, Polarized Light Microscopy, X-ray Diffraction, Scanning Electron Microscopy (SEM) & Transmission Electron Microscopy (TEM) leading some credibility to bacterial infection relationship

Bile is usually sterile in healthy individuals; however in presence of gallstones in gallbladder or biliary ducts, it could be colonized with bacteria. Detecting bactobilia could be valuable for planning antibiotic prophylaxis and treatment of postoperative infections after laparoscopic cholecystectomy. The overall incidence of bactobilia has been shown from 2.8% to 25% in different studies.

In this context, authors have confined themselves to studies of only pigment stones, leading to perceived high rate of

positive cultures. However, the data points to higher incidence of bacterial origin of Gallstones

I do acknowledge that a holistic data including all types of stones would have been more representative of the data and further research that is more inclusive needs to be carried out to find the true nature of causal relationship of stone & bile culture and gallstones

Dr. Rauf Ahmad Wani

MS; MRCS(ed); FACS

Professor, General & Minimal Invasive Surgery

SKIMS, Srinagar

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.768>

How to cite this article: Wani R. Bacterial Infection in etiology of Pigment Gallstones. JMS 2020; 23(3): 1

Received: 2020-08-05 Accepted: 2020-08-13

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Immune checkpoint inhibitor (ICPI) induced Nephrotoxicity An Insight

Mohmad Hussian Mir, Tariq Ahmad Bhat, Khalid Parvez Sofi, Imtiyaz Ahmad Wani, Muzafar Maqsood Wani

Department of Nephrology, SKIMS

There has been a marked increase in the therapeutic armamentarium of antineoplastic medications. Various classes of drugs acting at different sites have been developed over decades with the immune system being the latest target¹. The ability of the immune system to inhibit the cancerous growth of cells is called cancer immune-surveillance. Immune cells, primarily lymphocytes, recognize, and eliminate cells with spontaneous or inherited mutations that encourage malignant transformation^{1,2}.

The immune system acts in three phases to perform immune-surveillance. The **first phase** is the **elimination phase** in which the immune system eliminates transformed tumor cells before they become clinically evident. The **second phase** is the **equilibrium phase** where the tumor cells that survive the first phase are kept in check and prevented to proliferate further. The **third phase** is an **escape phase** in which tumor cells overcome immune defenses by loss of tumor antigens leading to a reduced recognition of tumor cells or by activation of immune checkpoint molecules such as cytotoxic T Lymphocyte antigen (CTLA-4) and programmed cell death protein 1 (PD-1). These checkpoints help keep immune responses from being too strong and sometimes can keep T cells from killing cancer cells¹.

Immune checkpoint inhibitors (ICPIs) may be defined as a drug that blocks checkpoints. When these checkpoints are blocked, T cells can better kill cancer cells. ICPIs resist cancer evasive mechanisms by allowing host immune cells to identify and eliminate malignant cells^{1,2}.

Pharmacologically, ICPIs are monoclonal antibodies targeting inhibitory receptors on T cells, tumor cells, and many other immune cells. Receptors including CTLA-4, PD-1, and PD-ligand 1 (PD-L1) are inhibited, activating T cells for antitumor activity^{1,2}. ICPIs currently in use are mentioned in Table 1. ICPIs are increasingly being used in patients with melanoma, non-small cell lung cancer, urothelial cancer, renal cell carcinoma and many other malignancies^{3,4}.

Table 1. Various classes of ICPIs available.

CLASS	DRUG
anti-CTLA-4 antibodies	Ipilimumab, Tremelimumab
anti- PD-1 antibodies	Nivolumab, Pembrolizumab, Cemiplimab, Spartalizumab
anti-PD L1 antibodies	Atezolizumab, Avelumab, Durvalumab, Pidilizumab.

Various side effects have been reported with ICPIs. In this review, we will be briefing about these side effects with emphasis on the nephrotoxic potential of ICPIs and their use in dialysis and renal transplant patients.

Immune-related adverse effects

Immune-related adverse events have been reported in >50%

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.774>

Correspondence:

Imtiyaz Ahmad Wani, DM
Professor, Department of Nephrology, SKIMS

How to cite this article: Mir MH, Bhat T, Sofi KP, Wani IA, Wani MM. Immune Checkpoint Inhibitor (ICPI) induced Nephrotoxicity An Insight. JMS 2020; 23(3): 2-5

Received: 2020-08-16 Accepted: 2020-08-24

of patients being treated with ICPIs with 20% of patients developing severe (grade 3) or life-threatening (grade 4) toxicities. Commonly affected organs are skin, gastrointestinal tract, liver, and endocrine organs. Commonly reported GI complications to include diarrhea, colitis, hepatitis, and constipation. Skin involvement includes rash, pruritus, eczema, sweet's syndrome, psoriasis Vulgaris, alopecia, and lichen planus. Endocrine side effects include hypophysitis, diabetes mellitus, pancreatitis, hypothyroidism, thyroiditis. Cardiac side effects reported are arrhythmia, pericarditis with effusion, cardiac tamponade, myocardial fibrosis, conduction blocks, troponin elevation, and myocarditis. Radiculitis, generalized epileptic seizures, aseptic meningitis, and myasthenia gravis have also been reported^{5,6}.

Nephrotoxicity

There are various forms of nephrotoxicity reported with ICPIs^{7,8}. From the asymptomatic rise in creatinine to active urinary sediment, proteinuria, hematuria, oliguria, all have been reported. Various kidney lesions have been documented in the setting of ICPIs therapy. The most common lesion described is acute interstitial nephritis (AIN), one-quarter of these lesions have associated granulomas⁹. Less commonly acute tubular necrosis (ATN), glomerulonephritis, and thrombotic microangiopathy has been reported^{10,11}.

Acute Kidney Injury (AKI)

First described in 2014, two patients on Ipilimumab, reported having granulomatous AIN¹¹. Six cases of AKI were reported in 2016 with biopsy-proven AIN following treatment with Nivolumab and Pembrolizumab¹². In another report, 13 patients developed AKI following exposure to ICPIs, AIN documented in 12, and granulomas in 3 of them. One patient had thrombotic microangiopathy. The overall incidence of ICPI-AKI is 2 to 3% based on data from various studies, reaching up to 5% in patients on combination ICPI therapy¹³. ICPI-AKI diagnosis is based on clinical features alone, and in some cases confirmed by kidney biopsy^{14,15}.

The National Cancer Institute defines AKI, by comparing changes in serum creatinine to the upper limit of normal. Kidney Diseases Improving Global Outcomes (KDIGO) defines AKI according to relative changes in AKI (Table 2). Another definition and classification system has been

proposed (Table 3)⁸. Severe AKI defined as an increase in serum creatinine more than threefold above baseline, an increase in serum creatinine to >4 mg/dl, or need of dialysis, was 0.6%¹³.

Table 2. Grading system for ICPI associated AKI

SYSTEM	GRADE	DESCRIPTION
CTCAE	Grade 1	Higher than ULN to 1.5× ULN
	Grade 2	> 1.5-3.0 × baseline; 1.5-3.0 × ULN
	Grade 3	3.0 × baseline; > 3.0 - 6.0 × ULN. Hospitalization indicated
	Grade 4	>6.0 × ULN; life -threatening consequences; RRT indicated
	Grade 5	Death
KDIGO	Stage 1	Increase in SCr of ≥0.3 mg/dl within 48 hrs or 1.5-1.9 × baseline
	Stage 2	Increase in SCr to 2-2.9 × baseline
	Stage3	Increase in SCr to 3 × baseline s or to ≥4.0mg/dl or initiation of RRT

CTCAE, Common Terminology Criteria for Adverse Events; ULN, Upper Limit of normal

Table 3. Definition and classification of ICPI-associated AKI

Classification	Definition
Definite ICPI-AKI	Kidney biopsy confirmed diagnosis compatible with ICPI-AKI and after clinical review of risk factors
Probable ICPI - AKI	BOTH of the following Sustained ↑ in SCr ≥50% on at least two consecutive values or need for RRT. Absence of an alternative plausible etiology AND at least one of the following Sterile pyuria (≥5 WBCs/hpf) Concomitant or recent extra renal irAE eosinophilia(≥500 cells/ul)
Possible ICPI-AKI	BOTH of the following ↑ in SCr ≥50% or need for RRT. AKI is not readily attributable to alternative causes

ICPI-AKI, immune checkpoint inhibitor-associated AKI, irAE, immune-related adverse event

Risk factors include prior or concomitant proton pump inhibitor use, prior or concomitant extra-renal irAEs.

Most patients present with rising creatinine, sterile pyuria, or sub nephrotic proteinuria. Eosinophilia is found in a minority of patients. After initiation of ICPIs median time to AKI is 14 weeks, much delayed in comparison to extra-renal side effects that appear within 4 weeks. Risk factors for developing AKI include concomitant use of proton pump inhibitors, combination treatment with anti-CTLA-4 and anti-PD-1/PD-L1 agents, and lower baseline GFR^{8,12,13}.

Patients who present with stage1 AKI should be evaluated for reversible causes like dehydration, hypercalcemia, urinary obstruction, or other nephrotoxic drug exposure.

Patients with persistent stage 1 and stage 2 or 3 AKI should undergo kidney biopsy. Empirical treatment with steroids should be considered only if there is a contraindication to performing kidney biopsy with an absence of alternative etiology for AKI⁸.

Glucocorticoids are the mainstay of treatment of ICPI induced tubulointerstitial nephritis, backed by observational data. The commonly used protocol is prednisolone 1mg/kg daily with a slow taper over 2-3 months. In patients with severe AKI, intravenous methylprednisolone 250-500 mg daily for 3 days may be used as initial therapy. Other agents used particularly in steroid-resistant patients include mycophenolate mofetil, cyclophosphamide, rituximab, and infliximab^{8,13}.

Usually, ICPI is withheld in patients with severe AKI. However, patients who have exhausted other options for chemotherapy may need to be rechallenged with ICPIs. In patients who develop grade 3 or higher AKI, ICPIs should be permanently discontinued according to the American Society of Clinical Oncology guidelines. But in one multicenter study thirty-one patients were rechallenged and only seven (23%) developed recurrent AKI, six of them showed partial or complete recovery. Rechallenge may be attempted in patients who have shown tumor response after a renal injury has resolved or stabilized⁸.

Glomerular disease. Nephrotic syndrome was first reported in a patient with metastatic melanoma, following ipilimumab treatment. Kidney biopsy revealed minimal change disease with complete foot process effacement. Complete resolution of proteinuria was achieved after discontinuation of ipilimumab and glucocorticoid treatment¹⁴. Collapsing Focal Segmental Glomerular Sclerosis (FSGS), with AKI has been reported following exposure to nivolumab, with no improvement after drug withdrawal and glucocorticoid treatment¹⁵. One patient with FSGS has been reported, with no improvement in drug withdrawal and glucocorticoid treatment, improved with mycophenolate mofetil¹⁶. Immune complex GN has been reported on exposure to ICPIs, including ipilimumab associated lupus-like glomerulopathy. Phospholipase A2 receptor antibody-negative membranous nephropathy has been reported in a patient on nivolumab, achieving partial recovery after the withdrawal of drug and glucocorticoid therapy¹⁷.

ICPIs in End-Stage Renal Disease (ESRD)

Experience of using ICPIs in ESRD is limited to case reports and small case series^{18,19}. A total of 26 patients, mostly on hemodialysis have been described in the literature²⁰. Ipilimumab, Nivolumab, Pembrolizumab, and Atezolizumab have been used in treating ESRD patients, for various malignancies including melanoma, renal cell carcinoma, squamous cell skin carcinoma, urothelial cancer, and non-small cell lung cancer. ICPIs are not renally excreted, so dose modification is not needed. Immune-related adverse events are similar to non-ESRD patients^{19,20}.

ICPIs in Kidney Transplant patients

Immunosuppressive drugs used in post-renal transplant patients predispose to cancer by inhibiting immune surveillance, suppressing DNA repair, and predisposing to oncogenic viral infections^{21,22}. Due to an increased predisposition to rejection with the use of ICPIs, transplant patients have been excluded from clinical trials of ICPIs. The use of ICPIs has been anecdotal^{23,24,25}. It has been observed that the rejection rate with Ipilimumab monotherapy was 33% compared to 52% with anti-PD-1 monotherapy and 55% with combination therapy²⁹. The strategies to prevent the high rate of rejection include the use of pre-emptive steroids and mammalian target of rapamycin (mTOR) inhibitors. Alternatively, calcineurin inhibitors may be continued during ICPI use. Depending upon the individual immunological risk, a decision about reduction or interruption of immunosuppression must be balanced with a high risk of rejection with ICPIs²⁶.

With ICPIs as the new class of antineoplastic medications, it is important that the oncologist, physicians, and the nephrologists be aware of various toxicities as well as its management.

REFERENCES

1. Leach DR, Krummel MF, Allison JP: Enhancement of antitumor immunity by CTLA-4 blockade. *Science* 1996; 271: 1734-1736.
2. Boussiotis VA. Molecular and biochemical aspects of the PD-1 checkpoint pathway. *N Engl J Med* 2016; 375: 1767-1778.
3. Callahan MK, Kluger H, Postow MA, Segal NH, Lesokhin A, Atkins MB, et al. Nivolumab plus ipilimumab in patients with advanced melanoma: updated survival, response, and safety data in a phase 1

- dose-escalation study. *J Clin Oncol* 2018; 36:391-398.
4. Garon EB, Rizvi NA, Hui R, Leigh N, Balmanoukian AS, Eder JP et al; KEYNOTE-001 Investigators: Pembrolizumab for the treatment of non-small-cell lung cancer. *N Engl J Med* 2015; 372: 20182028.
5. Postow MA, Sidlow R, Hellmann MD: Immune-related adverse events associated with immune checkpoint blockade. *N Engl J Med* 2018; 378: 158168
6. Sury K, Perazella MA, Shirali AC: Cardiorenal complications of immune checkpoint inhibitors. *Nat Rev Nephrol* 2018; 14: 571588.
7. Perazella MA, Shirali AC: Nephrotoxicity of cancer immunotherapies: Past, present and future. *J Am Soc Nephrol* 2018; 29: 20392052.
8. Shruti Gupta, Frank B. Cortazar, Leonardo V. Riella, David E. Leaf. Immune Checkpoint Inhibitor Nephrotoxicity: Update 2020. *Kidney* 2020; 360: 130-140
9. Shirali, A. C, Perazella, M. A. &Gettinger, S. Association of acute interstitial nephritis with programmed cell death 1 inhibitor therapy in lung cancer patients. *Am. J. Kidney Dis* 2016; 68:287291.
10. Cortazar, F. B. et al. Clinicopathological features of acute kidney injury associated with immune checkpoint inhibitors. *Kidney Int* 2016; 90, 638647.
11. Izzedine, H. Kidney injuries related to Ipilimumab. *Invest New Drugs* 2014; 32: 769773.
12. Izzedine H, Mathian A, Champiat S, Picard C, Mateus C, Routier E, et al : Renal toxicities associated with pembrolizumab. *Clin Kidney J* 2019; 12: 8188.
13. Seethapathy H, Zhao S, Chute DF, Zubiri L, Oppong Y, Strohbehn I et al: The incidence, causes, and risk factors of acute kidney injury in patients receiving immune checkpoint inhibitors. *Clin J Am Soc Nephrol* 14: 16921700, 2019
14. Kidd, J. M, Gizaw, A. B. Ipilimumab-associated minimal-change disease. *Kidney Int* 2016; 89, 720
15. Ray, A., Ghosh, S., Ghosh, M. &Yarlagadda, S. Nivolumab induced renal failure with collapsing focal segmental glomerulosclerosis (FSGS). *J. Am. Soc. Nephrol* 2016; 27, 1010A.
16. Daanen, R. A. et al. Nivolumab-associated nephrotic syndrome in a patient with renal cell carcinoma: a case report. *J. Immunother* 2017. 40, 345348.
17. Lin, J., Schiff, M., Salvatore, S., Shoustari, A. &Glezerman, I. Membranous nephropathy related to the checkpoint inhibitor nivolumab. *J. Am. Soc. Nephrol* 2016; 27, 102A.
18. Cheun H, Kim M, Lee H, OhK-H, Keam B. Safety and efficacy of immune checkpoint inhibitors for end-stage renal disease patients undergoing dialysis: A retrospective case series and literature review. *Invest New Drugs* 2019; 37: 579583.
19. Vitale MG, Baldessari C, Milella M, Buti S, Militello AM, Di Girolamo S et al. Immunotherapy in dialysis-dependent cancer patients: Our experience in patients with metastatic renal cell carcinoma and a review of the literature. *ClinGenitourin Cancer* 17: e903e908, 2019
20. Jamie S. Hirsch, RimdaWanchoo, Jia H. Ng, YurilyKhanin, and Kenar D. Jhaveri. Use of Immune Checkpoint Inhibitors in End Stage Kidney Disease Patients, Single Centre Experience and Review of Literature: *Kidney* 2020; 360 1:399-402.
21. Sprangers B, Nair V, Launay-Vacher V, Riella LV, Jhaveri KD: Risk factors associated with post-kidney transplant malignancies. *Clin Kidney J* 11: 315329, 2018
22. Hurkmans DP, Verhoeven JGHP, de Leur K, Boer K, Joosse A, Baan CC et al. Donor-derived cell-free DNA detects kidney transplant rejection during nivolumab treatment. *J Immunother Cancer* 2019; 7: 182.
23. Boils CL, Aljadir DN, Cantafio AW: Use of the PD-1 pathway inhibitor nivolumab in a renal transplant patient with malignancy. *Am J Transplant* 2016; 16: 24962497.
24. Spain L, Higgins R, Gopalakrishnan K, Turajlic S, Gore M, Larkin J: Acute renal allograft rejection after immune checkpoint inhibitor therapy for metastatic melanoma. *Ann Oncol* 2016; 27: 11351137.
25. Barnett R, Barta VS, Jhaveri KD: Preserved renal-allograft function and the PD-1 pathway inhibitor nivolumab. *N Engl J Med* 2017; 376: 191192.
26. Venkatachalam K, Malone AF, Heady B, Delos Santos R, Alhamad T: Poor outcomes with the use of checkpoint inhibitors in kidney transplant recipients. *Transplantation* 2020; 104(5):1041-1047

THE POTENTIAL IMPACTS OF OBESITY ON COVID-19

Tasaduk Sultan Khan¹, Musadiq Sultan Khan²

¹ Respiratory & General Medicine, Furness General Hospital, Barrow in Furness, Cumbria, UK

² Barnsley Hospital NHS Foundation Trust, Barnsley S75 2EP

Barrow in Furness, UHMBT, LA14 4LF, Cumbria, UK

Coronavirus disease 2019 is the outbreak caused by a new strain of coronavirus named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)¹. The outbreak that leads to a global pandemic originated from the city of Wuhan in China². According to a CDC report, the commonest underlying conditions among those hospitalized with COVID-19 were diabetes, chronic lung disease, and cardiovascular disease¹. Obesity is a common denominator for cardiovascular disease and diabetes. Previous experience with the H1N1 epidemic showed that obesity is associated with high mortality³. All of these observations have raised concerns about the impacts that obesity could have on COVID-19.

The impacts of obesity on respiratory physiology

In general, obesity is characterized by a decrease in total respiratory system compliance⁴. This is partially due to a decrease in lung compliance, which could be related to the increased pulmonary blood volume seen in obese subjects. The primary reason, however, is the decrease in chest wall compliance that is due to the accumulation of fat in and around the ribs and diaphragm and in the abdomen^{4,5}. To date, there are no published studies that mention the relation between obesity and the ventilator phenotypes of COVID-19⁶. However, an increased prevalence of obesity has been found among patients with SARS-CoV-2 pneumonia⁶.

The impacts of obesity on Renin- Angiotensin-Aldosterone System (RAAS)

Obesity is associated with an imbalance in the RAAS, resulting in an over-expression of the angiotensin II (ANG II) and angiotensin II receptor type 1 (AT1R) axis both at the level of adipose tissue and at systemic level⁷. It has been documented that the way adipose tissue acts as a reservoir for influenza A virus, HIV, human adenovirus, and cytomegalovirus, SARS-CoV-2 can also use the adipose tissue as a reservoir to affect other parts of the body⁸. Due to this obese subjects, especially those with extensive visceral adipose tissue, may develop an explosive systemic response of the ANG II and AT1R axis and could host and store an expansive viral load, which might lead to the development of a more severe form of the disease.

The impacts of obesity on immunity

Obese patients have a chronically lowered concentration of adiponectin (an anti-inflammatory adipokine) and higher levels of leptin (a pro-inflammatory adipokine)⁴. There is an excess production of leptin and other pro-inflammatory cytokines such as TNF alpha, IL-6, MCP-1, and IL-1β in adipose tissue, which leads to pro-inflammatory excess energy milieu. The abundance of these pro-inflammatory mediators in adipose tissue leads to dysfunction of the innate immunity under obesity^{4,5}.

Correspondence:

Dr. Tasaduk Sultan Khan, MD, MRCP(UK), FRCP(UK), FCCP(USA)
Consultant Respiratory & General Medicine,
Furness General Hospital,
Email: tasaduksultan.khan@mbht.nhs.uk

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.782>

How to cite this article: Khan TS, Khan MS. The Potential Impacts of Obesity on COVID-19. JMS 2020; 23(3): 6-7

Received: 2020-09-05 Accepted: 2020-09-07

Zhang et al hypothesized that leptin was a cofactor for the H1N1 pandemic in 2009⁹. The metabolic dysfunctions that are associated with obesity often lead to systemic leptin resistance. This resistance is associated with reduced bone marrow hematopoiesis and decreased T cell production and development in the thymus. Together, these disturbances result in diminished initiation of the immune responses, decreased memory T cell reactions, and decreased efficacy of vaccination^{4,5}. This impaired pro-inflammatory response has led to the severe lung lesions seen in patients of the H1N1 pandemic.

Obese patients are potentially more contagious than lean people

Studies have shown that obese patients are potentially more contagious than lean subjects in the setting of viral infections for the following possible reasons⁴.

1. Obesity increases the duration of viral shedding; symptomatic obese patients shed the virus 42% longer than people who do not have obesity.
2. The obesity state is characterized by a delayed capacity to produce interferons. This enables further viral RNA replication to increase the chances of new, more virulent viral strains emerging.
3. BMI was found to be positively associated with the level of infectious virus in the exhaled air.

Obesity in the clinical setting

Obese patients are more challenging to intubate, to diagnose via imaging, and for nursing staff to position and transport and may require special beds and positioning/transport facilities. From the ventilatory point of view, obese patients need high positive end-expiratory pressure (PEEP) and respond well to prone positioning and extracorporeal support⁴.

To conclude obesity has been found to be a strong independent risk factor for hospitalization in COVID-19 and to cause or exacerbate a variety of comorbidities which have been proven to be associated with increased morbidity and mortality among COVID-19 patients. Moreover, obese individuals have defective responses of both innate and adaptive immunity, with decreased response to vaccinations. And finally, obese patients are likely to strain the health system as they pose many challenges to healthcare provision inside and outside the ICUs.

REFERENCES

1. Centers for Disease Control and Prevention. 2019 *Novel Coronavirus, Wuhan, China*. CDC, 2020. Available from www.cdc.gov/coronavirus/2019-ncov/about/index.html
2. Zhou F, Yu T, Du R et al. Clinical Course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054-62.
3. Honce R, Cherry SS. Impact of obesity on Influenza A virus pathogenesis, immune response and evolution. *Front Immunol* 2019;10:1071
4. Kyrou I, Randeve HS, Tsigos C et al. Clinical problems caused by obesity. In; Feingold KR, Anawalt B, Boyce A et al (eds). *Endotext*. South Dartmouth(MA): MDText.com, Inc; 2000-2020.
5. Parameswaran K. Todd DC, Soth M. Altered respiratory physiology in obesity. *Can Respir J* 2006; 13:203-10.
6. Costa H, Jacob M, Pereira R, Calcas R. COVID 19 ventilatory phenotypes, and obesity: is there a relationship? *Obesity* 2020, in press (doi:10.1002/oby.22877).
7. Pinheiro TA, Barcala-Jorge AS, Andrade JMO et al. Obesity and malnutrition similarly alter renin-angiotensin system and inflammation in mice and human adipose. *J Nutr Biochem* 2017;48:74-82.
8. Bourgeois C, Gorwood J, BBarrail-Tran A et al. Specific biological features of adipose tissue, and their impact on HIV persistence. *Frontiers Microbiol* 2019;10:2837.
9. Zhang AJX, To KKW, Lau CCY, et al. Leptin mediates the pathogenesis of severe 2009 pandemic influenza A (H1N1) infection associated with cytokine dysregulation in mice with diet-induced obesity. *J Infect Dis* 2013;207: 1270-80

Study of Bacterial Aetiogenicity of Pigment Gall Stones by Culture of Nidus of Calculus and correlation between Stone Culture and Bile Culture

Mahmood Rashid Shiekh¹, Tahleel Shera², Shiekh Aamir Rashid³, Altaf Hussain Shera⁴, Junaid Ahmad⁵

^{1,4} Department of General Surgery, Govt. Medical College, Baramulla

² Department of Radio Diagnosis, Govt. Medical College, Baramulla

³ ASCOMS, Jammu

⁵ Department of Microbiology, Govt. Medical College, Baramulla

ABSTRACT

Background: Calculus disease of biliary tract is a major public health problem in north Kashmir and pigment gall stones form a significant portion of gall stones in our region. In this study we prospectively studied 84 cases of pigment gall stones to examine the role of bacteria in the formation of these stones.

Objective: To study the bacterial aetiology of pigment gall stones.

Methods: A hospital based prospective study of 84 consecutive cases of pigment gall stones was done from December 2012 to November 2019 and the results of stone culture and bile culture were statistically analysed. Informed consent was obtained from all the patients in the study group.

Results: A total of 84 cases studied included 32 (38.09%) male patients and 52 (61.91%) female patients. The median age was 48.5 years and mean age was 46.2 years. Of 84 patients, 21 (25.00%) had black pigment stones and 63 (75.00%) had brown pigment stones. 61 (72.62%) patients presented with biliary colic and 24 (28.57%) patients had chronic cholecystitis. Bile culture was positive in 46 (54.76%) patients and stone culture was positive in 70 (83.33%) patients.

Conclusion: A strong association was observed between stone culture and bile culture results (showing similar beta-glucuronidase producing bacteria) and pigment stone formation.

Key words: Gall stones, stone culture, bile culture, nidus

JMS: 2020; 23(3): DOI: <https://doi.org/10.33883/jms.v23i3.746>

INTRODUCTION

Gall stone disease is known to mankind since ancient times.¹ Pigment gall stones have been referred to as early as 500 AD by Alexander of Tralle². These constitute 10% to 27% of all types of gall stones found in general population and both intrahepatic gall stones and stones caused by bile stasis or cirrhosis are usually pigment stones³. Pigment stones are usually multiple, 2 to 5mm in diameter, irregular to smooth in shape, black to brown in colour and amorphous or crystalline on cross section. Pigment stones, especially the brown ones, can form within extra hepatic and intra hepatic ducts apart from their usual occurrence in the gall bladder

and are commonly associated with ascending cholangitis. Enteric bacteria have been found to play a significant role in pigment gall stone formation and these stones in general and brown ones in particular have a tendency to recur while black gall stone disease is usually non recurrent⁴. Unlike the causative factors like cirrhosis, ileal resection etc. observed in the West, infection has been found to be a predominant cause in South East Asia.¹⁰ In this study, we prospectively

Correspondence:

Dr. Shiekh Mahmood Rashid
Assistant Professor, General Surgery,
Govt. Medical College, Baramulla
Email: mahmoodshiekh234@gmail.com

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.746>

How to cite this article: Anand S, Ahmed O, Kundal V. Clinical spectrum and outcomes of snake bite patients admitted in a tertiary care hospital, A prospective observational hospital based study. JMS 2020; 23(3):8-12

Received: 2020-05-09 Accepted: 2020-08-11

studied 84 patients with pigment gall stones disease to examine the role of beta glucuronidase producing bacteria in the formation of these type of stones. 'Pigment gall stones' included any black or dark brown stone containing less than 25% cholesterol, whether or not pigment was its major component and irrespective of the nature and proportion of other constituents¹³.

MATERIAL AND METHODS

A hospital based prospective study was carried out from December 2012 to November 2019 in the departments of General Surgery, Microbiology and Radio diagnosis of Govt. Medical College Baramulla, Kashmir. It included a total of 84 consecutive patients with pigment gall stones (confirmed by chemical analysis of extracted gall stones). A formal informed consent was taken from all of them. The patients were selected as they fulfilled the following inclusion criteria: confirmed to have pigment gall stones, that is, black to dark brown gall stones containing less than 25% cholesterol irrespective of the proportion of other constituents and whether are not pigment was its major constituent²⁰; not on any medication for gall stones for last three months and had not received antibiotics for two weeks prior to surgery.

A detailed account of history was taken and complete physical examination was performed. Routine investigations including complete blood count, liver function tests, serum amylase and calcium levels were requested as also an abdominal ultrasound performed to know about the location, number and size of gall stones and gall bladder wall thickness or oedema and any associated dilatation of hepato-pancreatic ducts was ruled out. In selected cases, an abdominal computerised axial tomogram or a magnetic resonance cholangiogram was also done to exclude or to elaborate on any added pathology.

All the 84 patients were subjected to cholecystectomy and gall bladder sample along with the stones within it was placed aseptically in a container. 5 to 10 ml of bile was aspirated from within the gall bladder and preserved for bile culture and sensitivity in all the 84 cases during the operative procedure. The gall stones were removed and immediately transferred to a McCartney bottle containing 5ml of normal saline for further processing. A portion of stone was sent for chemical analysis and the other portion preserved for microbiological evaluation. Only those

calculi were included in the present study whose chemical analysis results fulfilled the criteria already stated for the case of pigment stones.

A technique for culture of calculi as devised by Nemoy and Stamey²² in 1971 was done. The culture methods were designed for localization of bacteria to the substance of the calculus and to distinguish bacteria within a stone from surface colonization. Under all aseptic precautions, a portion of stone was washed consecutively in four test tubes containing 5 ml of sterile saline each and then crushed in another tube in a laminar air flow chamber. 0.1ml of fluid from washes 1 and 4 and of the fluid of crushed stones were quantitatively cultured in Blood Agar, McConkay Agar and Brain Heart Infusion Broth and Agar. The cultures were studied at 24 hours and 48 hours and organisms typed. An increased density of isolate in the fluid of crushed stone and increased colony count after culture compared with the fluid of wash 4 was considered evidence for bacteria within a stone. The spectrum of bacteria that colonized the pigment stone was detailed out to assess their relation with the genesis of pigment gall stones.

RESULTS

In the present study of 84 consecutive patients with pigment gall stones, confirmed by stone analysis, conducted in the department of General Surgery at Govt. Medical College Baramulla, 32(38.09%) patients were males and 52(61.91%) patients were females. The median age was 48.5 years and mean age was 46.2 years. The distribution of study subjects as per the type of stones found in 84 patients showed that 21(25.00%) patients had black pigment stones and 63(75.00%) patients had brown pigment gall stones. (Table 1).

Table 1: Distribution of 84 study subjects as per type of stone.

Type of pigment stone	Number	Percentage (%)
Black pigment stones	21	25.00
Brown pigment stones	63	75.00
Total	84	100

Of 84 study subjects, 61 (72.62%) patients presented with biliary colic, and 24(28.57%) patients presented with chronic cholecystitis. Acute infective conditions like acute cholecystitis were observed in 2(2.38%) patients. 16(19.05%) patients presented with ultrasonic evidence of mucocele or empyema of gall bladder prior to surgery. Only 1(1.19%) patient had a documented predisposing cause in

the form of glucose 6 phosphate deficiency haemolytic anaemia. (Table 2).

Table 2: Distribution of cases as per the presenting features .

Presenting feature	Number	Percentage (%)
Biliary colic	61	72.62
Chronic cholecystitis	24	28.57
Acute cholecystitis	02	2.38
USG evidence of infection	16	19.05
Predisposing causes in the form of G6PD deficiency haemolytic anaemia	01	1.19

Intra operative findings revealed thickened gallbladder wall suggesting chronic cholecystitis in 46 (54.76%) patients; 3(3.57%) patients had dense adhesions of gallbladder with surrounding viscera and 3 (3.57%) patients had gall bladder perforation with cholecystoenteric fistula formation. (Table 3).

Table 3: Presentation of subjects at the time of surgery.

Intraoperative presentation	Number	Percentage (%)
Thickened gallbladder wall	46	54.76
Gallbladder perforation	03	3.57
Adhesions	03	3.57

Of 84 study subjects, bile culture was positive in 46 (54.76%) cases while stone culture was positive in 70 (83.33%) cases. Both bile and stone cultures were positive in 32 (38.09%) cases; bile culture was negative but stone culture was positive in 38(45.24%) cases while in 14 (16.67%) cases both the cultures were negative.(Table 4). The stones which were positive showed uniform distribution of bacteria throughout the stone including its core as confirmed by the culture and sensitivity reports of the fragments of the stone including its core.

Table 4: Comparison of stone culture and bile culture results of 84 study subjects

Culture positivity	Number	Percentage (%)
Bile culture positive	46	54.76
Stone culture positive and bile culture positive	32	38.09
Stone culture positive and bile culture negative	38	45.24
Stone culture negative	14	16.67
Bile culture negative	38	45.24

Of 70(83.33%) culture positive stone culture results, 11(13.09%) black stones and 59(70.24%) brown stones were culture positive for bacteria (Table 5).

Table 5: Stone culture results of 84 study subjects.

Evidence of bacteria	Number	Percentage (%)
Stone culture positive	70	83.33
Stone culture negative	14	16.67
Presence of bacteria in black stones	11	13.09
Presence of bacteria in brown stones	59	70.24

Most of the bacteria were bacillary forms and *E. coli* was the commonest organism grown in 73.81% of stone cultures. Few cultures revealed other types of bacteria such as *Klebsiella* (3.57%), *Salmonella typhi* (3.57%) and *Staphylococcus aureus*. (Table 6). In some of the samples, there were groups of bacteria suggesting colony formation.

Table 6: Distribution of 70 patients with culture positive stone nuclei in relation to bile culture results.

Bacteria	Number of patients with culture positive stone cultures		Bile culture showing same bacteria	
	Number	percentage (%)	Number	Percentage (%)
<i>E.coli</i>	62	73.8	28	33.33
<i>Klebsiella</i>	3	3.57	1	1.19
<i>Salmonella Typhi</i>	3	3.57	2	2.38
<i>Staphylococcus aureus</i>	2	2.38	1	1.19
Total	70	83.33	32	38.09

DISCUSSION

In the present study, out of 84 patients studied, 32(38.09%) patients were males and 52 (61.91%) patients were females. The median age was 48.5 years and mean age was 46.2 years. Of 84 patients studied, 21(25.00%) patients had black pigment stones and 63(75.00%) patients had brown pigment stones. 72.62% patients presented with biliary colic and 28.57% patients presented with chronic cholecystitis. At surgery 54.76 % patients had thickened gallbladder wall suggesting chronic inflammation and 3.57 % cases presented with gallbladder perforation. These findings correlated with those of Akiyoshi T and Nakayama F²⁰. The central portions (nidus) of the stones collected from 84 patients were cultured as per the technique already described. It was observed that 83.33% of the stones from 70 patients were culture positive and 16.67% of the stones from 14 patients were cultured negative. Of 83.33 % culture positive patients, beta glucuronidase producing organisms were found in 73.81 % of patients. Of 70 culture positive stone cases, 32(38.09%) had bactobilia with same organisms. These findings suggest that bacterial infection with beta glucuronidase producing organisms plays a significant role in pigment gallstone formation and pigment gall stones are the tomb stones of bacteria; in addition it was observed that majority of black pigment stones were not associated with predisposing causes such as ileal resection, cirrhosis of liver or haemolytic disease unlike the observations in Western studies²¹ and were again found to be associated with infection of stone nuclei and bile. Authors

wanted to explore at this aspect and especially to explore the association of black pigment stones with bacterial infection of bile and stone material. It was found that a significant portion (25.00%) of our study subjects who had black pigment stones had no predisposing factors mentioned above in Western world and were again found to be associated with recurrent biliary infection.

Though it was observed that bacteria are possibly etiological agents due to the evidence at hand, we cannot exclude the possibility that they are innocent bystanders. The fact that bacteria were found in the nuclei of multi-layered stones favours the hypothesis that bacteria are associated with stones at a nascent stage of their development. These findings are supported by the work of Soloway RD et al^{24,25}. The possible mechanism in pigment stone formation could be a combination of cholecystitis and bile duct occlusion due to bacterial infection producing excessive mucous formation and release of deconjugated bilirubin, bile acids, hydrolyzing phospholipids and calcium, thereby leading to pigment stone formation^{5-9, 11, 12}. Similar observations have substantiated the close association of bacteria with brown pigment stone formation^{15-19, 14, 23}.

CONCLUSION

Considering all these findings, it could be suggested that hepatobiliary tract infection with beta glucuronidase producing bacteria has a significant role in the formation of pigment gall stones. Bacteriological study of a large number of stones especially in relation to their age, chemical composition and serial culture of patients might help in further establishing the role of bacterial infection in stone formation.

Financial support and sponsorship: Nil

Conflict of Interest: There are no conflict of interest

REFERENCES

1. Berci G. Historical overview of surgical treatment of biliary stone disease. In: MacFadyen BV, Arregui M, Eubanks S, Olsen DO, Peters JH, Soper NJ, et al., *Laparoscopic Surgery of the Abdomen*. New York: Springer; 2004: 139-142.
2. Alexander of Tralle, cited by Rains AJH: Gall stones: Causes and Treatment. Springfield, Ill, Charles C Thomas, 1964, p.1.
3. Baig SJ, Biswas S, Das S, Basn K, Chatopadday G. Histopathological changes in gall bladder mucosa in cholelithiasis: Correlation with chemical composition of gall stones. *Trop Gastroenterol*. 2002; 23: 25-27.
4. Branganza JM, Worthington H.A. Radical view of gall stone aetogenesis. *Med Hypotheses*. 1995; 45: 510-516.
5. Shiesh SC, Chen CY, Lin XZ, Lin ZA. Melatonin prevents pigment gall stone formation induced by bile duct ligation in guinea pigs. *Hepatology*. 2000; 32: 455-460.
6. Lin XT, Hu J. Relationship between bilirubin free radical and formation of pigment gall stones. *World J Gastroenterol*. 2002; 8: 413-417.
7. Haigh WG, Lee SP. Identification of oxysterols in human bile and pigment gall stones. *Gastroenterology* 2001; 121: 118-123.
8. Leonard MR, Andrade JD. Striking alterations in bilirubin and bile salt solution chemistry of gall bladder bile in a germ free mouse model of black pigment gall stones. *Gastroenterology*. 2011; 140 suppl 1: 568.
9. Whary MT, Woods SE, Fox JG. Pigment gall stones in germ free Swiss Webster mice. *Gastroenterology*. 2010; 138 suppl 1: S211.
10. Fan Y, Wu SD, Sun L. Possible relationship between intestinal barrier function and formation of pigment gall stones in hamsters. *Hepatobiliary Pancreat Dis Int*. 2008; 7: 529-532.
11. Rahman GA. Cholelithiasis and Cholecystitis: changing prevalence in an African community. *J Natl Med Assoc*. 2005; 97: 1531-1538.
12. Ostrow JD. Unconjugated bilirubin and cholesterol gall stone formation. *Hepatology*. 1990; 12: 219S-226S.
13. Donovan JM, Carey MC. Physicochemical basis of gall stone formation. *Gastroenterol Clin North Am*. 1991; 20: 47-66.
14. Ostrow JD, Mukerjee P. Revalidation and rationale for high pKa values of unconjugated bilirubin. *BMC Biochem*. 2007; 8: 7.
15. Vang L, Shen W, Wen J. An animal model of black

pigment gall stones caused by nanobacteria. *Dig Dis Sci.* 2006; 51:1126-1132.

16. Vitak L, Carey MC. Enterohepatic cycling of bilirubin as a cause of black pigment gall stones in adult life. *Eur J Clin Invest.* 2003; 33: 799-810.
17. Aydogdu I, Sari R, Ulu R, Sevince A. The frequency of gall bladder stones in patients with pernicious anaemia. *J Surg Res.* 2001; 101: 120-123.
18. Diehl AK, Holleman DR. Clinical correlates of gall stone composition: distinguishing pigment from cholesterol stones. *Am J Gastroenterol.* 1995; 90: 967-972.
19. Hoffman AF, Ton Nu HT, Mysels KJ. Solubility of calcium salts of unconjugated and conjugated natural bile acids. *J Lipid Res.* 1992;33:635-646.
20. Akiyoshi T, Nakayama F. Bile acid composition in brown pigment stones. *Dig Dis Sci.* 1990;35:27-32.
21. Kaufman HS, Magnuson TH, Lillemoe KD. The role of bacteria in gall bladder and common duct stone formation. *Ann. Surg.* 1989; 209: 584-591.
22. Nemoy NJ. Surgical, bacteriological and biochemical management of infection stones. *JAMA.* 1971;215: 1470-76.
23. Stewart L, Oesterle AL, Erdan I, Griffiss JM, Way LW. Pathogenesis of pigment gall stones in Western societies; the central role of bacteria. *J Gastrointest Surg.* 2002; 6(6): 891-930.
24. Soloway RD, Trotman BW, Ostrow JD, Pigment versus Cholesterol cholelithiasis: comparison of stone and bile composition. *Am J Dig Dis.* 1974;19:585-90.
25. Trotman BW, Soloway RD. Pigment versus Cholesterol cholelithiasis; clinical and epidemiological aspects. *Am J Dig Dis.* 1975;20:735-40.

Clinical spectrum and outcomes of snakebite patients admitted in a tertiary care hospital. A prospective observational hospital-based study

Sameer Anand¹, Owais Ahmed Wani², Vijay Kundal³

^{1,3} Department of General Medicine, GMC Jammu.

² Department of Cardiology GMC Jammu.

ABSTRACT

Background: Snakebite is a major public health problem and its prevalence is high in India. Insufficient epidemiological data and global neglect of this condition prompted the World Health organization to recognize it as a "neglected tropical disease"

Objective: To study demographic characteristics of snakebite victims, to see clinical symptoms of bite and outcomes of snakebite.

Methods: A prospective observational hospital-based study carried out from November 2016 to October 2017, in patients with a history of snakebite.

Results: Total 300 patients were included in this study, 170 (56.67%) were bitten by hemotoxic snakes and 130 (43.33%) were bitten by neurotoxic snakes. Majority of patients were in the age group of 20-39 years (n=153; 51%). Men outnumbered women (n=190; 63.33%), with male to female ratio of 1.7: 1. Higher incidence of snake bite was found in July-September (n=135; 45.00%) followed by April to June (n=124; 41.33%). Majority of patients were farmers in rural areas (n=242; 80.80%). Out of 300 patients, 80 required ICU care. In patients with neurotoxic snake bites, clinical features were: ptosis (n=126; 96.92%), ophthalmoplegia (n=98; 75.38%), respiratory paralysis (n=60; 46.15%), bulbar weakness (n=74; 56.92%) and abdominal pain (n=25; 19.23%). Clinical features in patients of hemotoxic snake bites were: bleeding from bite site (n=110; 65.29%), cellulitis (n=100; 58.82%), gum bleed (n=53; 31.18%), ecchymosis (n=49; 28.82%), epistaxis (n=27; 15.82%), gastrointestinal bleeding (n=25; 14.71%) and haemoptysis (n=19; 11.18%). Complications observed were acute kidney injury (n=62; 20.67%), respiratory failure (n=51; 17.00%), DIC (n=7; 2.33%) and ARDS (n=5; 1.67%). Out of 170 hemotoxic snake bite patients, 157 (92.35%) patients recovered and 13 (7.65%) patients died and out of 130 neurotoxic snake bites, 124 (95.39%) patients recovered and 6 (4.61%) patients died, overall mortality was 6.3%. In our study, bite to needle time was less than 1 hour in 47 patients (15.66%), less than 6 hrs in 120 patients (40%) and more than 24 hours in 45 (15%).

Conclusion: Snakebite is common in adult males between 20 to 50 years and the commonest site is the lower limb. The majority of the victims are farmers who work in fields. A maximum number of cases presented within 1-6 hours of bite using a tourniquet on the affected limb as first aid. The complications and mortality of snakebite are high.

Keywords: Neglected tropical disease, Indian cobra (Naja naja), common krait, snakebite

JMS; 2020; 23(3) p?-p? DOI: <https://doi.org/10.33883/jms.v23i3.751>

INTRODUCTION

Snakebite counts significant health burden in the South East Asian Region resulting in death and/or disability of young individuals involved in agriculture work. Approximately 2,000,000 snake bites are reported in India each year. The million death study revealed 45,900 national snakebite deaths comprising 5% of all injury deaths and 0.5% of all deaths in India.¹ However there is no exact figure of mortality and morbidity associated with snakebite due to

poor reporting in the region.² Among 250 species of snakes found in the South East Asian Region, only 60 species are venomous included in three families, Elapidae, Viperidae and Colubridae.³ Around four million snake bites occur each

Correspondence:

Dr. Owais Ahmed Wani, DNB
Senior Resident, Department of Cardiology GMC Jammu.
Email: docowaiss@gmail.com

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.751>

How to cite this article: Anand S, Ahmed O, Kundal V. Clinical spectrum and outcomes of snake bite patients admitted in a tertiary care hospital, A prospective observational hospital based study. JMS 2020; 23(3):13-17

Received: 2020-05-27 Accepted: 2020-08-29

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

year in Asia, resulting in 100,000 deaths per year. In South Asia, four species are responsible for most of the deadly cases of venomous snake bites. These have been referred to as the 'big four', and include the Indian cobra (*Naja naja*), the common krait (*Bungarus caeruleus*), Russell's viper (*Daboia russelii*), and the saw-scaled or carpet viper (*Echiscarinatus*)⁴. Jammu region, which lies in the sub-Himalayan region is enormously infested with various types of poisonous snakes belonging to families Elapidae and Viperidae. Those belonging to viperidae are mainly hemotoxic and the commonest in Jammu province⁵. *Echiscarinatus* (saw-scaled viper) and Russell viper account for more than 85% of poisonous bites in the Jammu region⁶. The most notorious venomous snakes can be recognized by their size, shape, color, the pattern of markings, behavior, and the sound they make when they feel threatened. Viperidae (all species) results in local envenoming (swelling) with bleeding/clotting disturbances. Russell's viper cause local envenoming with shock or acute kidney injury. King cobra causes local envenoming with paralysis while Krait cause paralysis with minimal or no local envenoming². Though snake bite is a common health problem, there is no exact nationwide data available in India. This study aimed to see the demographic characters of the snakebite victims at Jammu city and other factors like site of the bite, type of bite, envenoming signs, delay of hospital presentation, first aid measure before hospital presentation, mortality, and morbidity.

MATERIAL AND METHODS

This was a prospective observational study carried out from November 2016 to October 2017. Inclusion criteria were, all the adult patients of poisonous snake bite admitted in medical wards and intensive care units. Exclusion criteria were all patients of non-poisonous snake bite & scorpion bite. The study was done at the Postgraduate Department of General Medicine, GMC Jammu. After obtaining permission from the ethical committee, the records of snake bite and suspected snakebite cases presenting to ER over a period of 1 year were collected and analyzed for demographic characteristic, site of bite; time elapsed since the bite, pre-hospital care, bite marks, presenting symptoms, clinical signs of envenoming and condition at discharge. Records of collected data were analyzed for the final outcome. Standard statistical methods were used in the

analysis of the data using Microsoft Excel.

RESULTS

Total 300 patients were enrolled in this study. There were 190 males and 110 females, with a male: female ratio 1.7:1. Majority (n=82; 27.33%) of patients were in the age group of 20-29 years followed by 30-39 years age group (n=71; 23.67%) [Figure 1]. Majority of patients were farmers (n=242; 80.80%). Next in the list were housewives (n=28; 9.20%). Majority of patients (n=203; 67.67%) were from rural areas. Higher incidence of snake bite was found in July-September (n=135; 45.00%) followed by April to June (n=124; 41.33%). 255 (85%) patients presented to emergency within 24 hrs, 45 (15%) reached after 24 hours, 47 (15.67%) reached within 1 hour while as 120 (40%) reached within 1-6 hours.

Fig 1. Distribution of patients in various age groups.

The majority of patients (n=170; 56.67%) were bitten by hemotoxic snakes and 130 (43.33%) patients were bitten by neuroparalytic snakes. In this study, the majority (n=200; 66.67%) of patients were bitten on lower extremity followed by upper extremity (n=67; 22.33%) and only 33 (11%) patients were bitten on head and trunk region. In the lower extremity, out of 200 cases, the majority (n=151; 88.82%) were hemotoxic snake bites. In the upper extremity, out of 67 cases, the majority (n=48; 36.92%) were neurotoxic snake bites. In the head and trunk region, all (n=33; 11.00%) cases were neurotoxic snake bites.

Prehospital treatment was very common (n=245; 81.67%). Among that use of tourniquet on the affected limb was the highest (n=151; 51.33%) followed by wound incision (13.6%), application of herbs (10.33%) and snake stone (6.33%) on the affected area, 55 (18.3%) patients did not receive any first aid prior to hospitalization. Among the total patients, 131 (43.67%) had local pain, tenderness, swelling and local rise of temperature, 56 patients (18.67%) had blisters, 40 patients (13.3%) had discoloration,

tingling, and burning sensation, 19(6.3%) patients had ulceration. Majority of patients (n=201; 67%) had double fang marks, 53 (17.6%) had single fang marks and only scratches were present on 18 (6 %) patients. Out of 300 patients, 80 patients with respiratory paralysis and ARDS required ICU care.

Clinical features in patients with neuroparalytic snake bites are shown in Figure 2, while clinical features in patients with haemotoxic snake bites are shown in Figure 3.

Fig 2. Clinical features in neuroparalytic snake bites patients.

Ptosis(n=126; 96.92%), ophthalmoplegia (n=98; 75.38%), respiratory paralysis (n=60; 46.15%), bulbar weakness (n=74; 56.92%), abdomen pain (n=25; 19.23%) and loss of consciousness (n=19; 16.15%)

Fig 3. Clinical features in haemotoxic snake bites patients.

Bleeding from bite Site (n=110; 65.29%), cellulitis (n=100; 58.82%), gum bleed (n=53; 31.18%), ecchymosis (n=49; 28.82%), epistaxis (n=27; 15.82%), gastrointestinal bleeding (n=25; 14.71%), haemoptysis (n=19; 11.18%) and intracranial bleeding (n=3; 1.76%)

Complications observed were acute kidney injury (62; 20.67%), respiratory failure (51; 17.00%), gangrene at the bite area (10; 3.33%), DIC (2.33%), ARDS (1.67%) and

intracerebral haemorrhage (1.00%). In the present study, the majority of patients (n= 170; 56.67%) had 3-7 days of hospital stay, 76 (25.33%) had a hospital stay of <3 days and 54 (18.00%) had a hospital stay of >7 days. Out of 170 haemotoxic snake bites, 157 (92.35%) patients recovered and 13 (7.65%) patients died. Out of 130 neurotoxic snake bites, 124 (95.39%) patients recovered and 6 (4.61%) patients died, overall mortality was 6.3%. In our study, bite to needle time was less than 1 hour in 47 (15.6%) patients, less than 6 hrs in 120 (40%) patients and more than 24 hours in 45. The majority of patients (40 %) presented to an emergency within 1-6 hrs of bite followed by 29.33% in 6-24 hrs, 15 % of patients presented after 24 hours.

DISCUSSION

In India, there is yet no large nationwide data bank to show the exact snakebite incidence and its sequelae i.e. morbidity and mortality. WHO has recommended that snake-bite should be formally recognized as an important occupational disease in the South East Asian region. And it is shown in studies that in South Asia, Southeast Asia, sub-Saharan Africa exists the highest burden of morbidity and mortality due to snake bites⁷. Despite its importance, there have been fewer proper clinical studies and there is a lack of reliable epidemiological data nationwide.

In this study, most of the snakebite patients were young males (63.33%) which is in corroboration with previous studies⁸. Individuals in this age group are actively involved in farming & outdoor activity thus vulnerable to this occupational hazard. The commonest site of the bite was in the lower limb in our study (n=200) and finding correlates with the previous literature^{3,7}. The mortality rate was around 6% in our study, In a North Indian study by Sharma et al mortality rate was 3%⁹. The complications encountered in our study were bleeding from bite site (65.29%), cellulitis (58.82%) and AKI (20.67%). In their study Bhat RN et al.⁶ and Saini et al⁹ recorded bleeding tendency in 65% and 47.80% respectively.

In a study by Nagnath R et al. in Central India, most vasculotoxic bites were associated with cellulitis (90.60%), coagulopathy (19%), and respiratory paralysis was encountered in 19.20% of the cases¹⁰. In our study, respiratory paralysis was noted in 60 (20%) patients and bulbar weakness in 74 (24.66%) patients, 10% of the cases required mechanical ventilatory support. The incidence of

AKI was 14.4% and 43.3% in studies by Halesha BR et al¹¹. and Nagnath R et al¹⁰ respectively. These differences in the complications found in different studies imply differences between the venoms of the subspecies unique to that geographic area. The importance of bite to needle time is a significant factor in influencing mortality. A study from Nepal showed that the delay in receiving treatment was significantly longer for victims with a fatal outcome¹². The optimum time for administration of ASV has been previously suggested as “as soon as possible” or within 6 hours. Correlation between the door to needle time and mortality is similar to those observed by Kavitha Saravu et al¹³. and I F Inamdaret al¹⁴. In the study by Nagrath et al¹⁰, mortality was high (66.7%) in patients with a bite to needle time more than 24 hours, as compared to patients whose bite to needle time was less than 24 hours and lowest when the bite to needle time was less than 6 hrs (7.2%). In our study, bite to needle time was less than 1 hour in 47 patients and less than 6 hrs in 120 patients, more than 24 hours in 45 patients and overall mortality was 6.33 %

The chief limitation of our study is that it is based on a prospective analysis of a small number of cases (n = 300). It is imperative to carry out a larger clinical study to better understand about clinical spectrum and outcomes.

CONCLUSION

Snakebite is a common medical emergency, where timely treatment can reduce morbidity and mortality and save precious human lives. Lack of information about simple measures of prevention, occupational hazard risks, and inappropriate first-aid measures all magnify the risk. Poor access to health care services, difficult transportation, and consequent delay in ASV administration results in a high fatality. The time elapsed after the bite is of vital importance because with the passage of time more venom gets bound to the tissues and is thus less manageable for neutralization by ASV. Many bites could be avoided by educating the population at risk. Sleeping on a cot (rather than on the floor) and under bed nets decreases the risk of nocturnal bites. Many bites occur when people walking barefoot or wearing only sandals accidentally step on a snake. Using a torch/ flashlight while walking on footpaths at night, and wearing boots and long trousers during agricultural activities, could significantly reduce the incidence of bites. Thus, we need an organized effort to cope with important aspects of snakebite management; foremost being, to

educate the rural population, first-aid management of the patient, and prompt medical management of emergencies, especially in rural areas.

REFERENCES

1. Mohapatra B, Warrel DA, Suraweera W, Bhatia P, Dhingra N *et al.* (2011) Snakebite Mortality in India: a nationally Representative Mortality Survey. *Plo S Negl Trop Dis* 5 (4): e1018. doi: 10.1371/journal.pntd.0001018.
2. Warrell DA. Epidemiology of snake-bite in the South-East Asia Region. In: Warrell DA (ed.) Guidelines for the management of snakebite. New Delhi WHO Press: New Delhi, India: WHO regional office for Southeast Asia. 2010;1-135.
3. Hansdak SG, Lallar KS, Pokharel P, Shyangwa P, Karki P, Koirala S. A clinic epidemiological study of snake bite in Nepal. *Trop Doct.* 1998 Oct; 28(4): 223-6. <https://www.ncbi.nlm.nih.gov/pubmed/9803844>.
4. Alirol E, Sharma SK, Bawaskar HS, Kuch U, Chappuis F. Snake bite in South Asia: a review. *PLoS Negl Trop Dis.* 2010 Jan; 4(1): e603. doi: 10.1371/journal.pntd.0000603 PMID: PMC2811174.
5. Kapoor S. Spectrum of snake bite poisoning in Jammu- an observational study. *J. Evid. Based Med. Healthc.* 2017; 4(88), 5139-5146. DOI: 10.18410/jebmh/2017/1027.
6. Bhat RN. Viperine snake bite poisoning in Jammu. *J Indian Med Assoc.* 1974;63(12):383-392.
7. Kasturiratne A, Wickremasinghe AR, De Silva N, Gunawardena NK, Pathmeswaran A, Premaratna R, et al. The Global Burden of snakebite: A Literature Analysis and Modelling Based on Regional Estimates of Envenoming and Deaths. Winkel K, editor. *PLoS Med* [Internet]. 2008 Nov 4; 5(11): e218. Available from: <http://dx.plos.org/10.1371/journal.pmed.005021>
8. Anjum A, Husain M, Hanif SA *et al.* (2012) Epidemiological Profile of Snake Bite at Tertiary Care Hospital, North India. *J Forensic Res* 3: 146. doi: 10.4172/2157-7145.1000146
9. Sharma N, Chauhan S, Faruqi S *et al* (2005) Snake envenomation in a north Indian hospital.
10. Saini RK, Arya RK, Singh *et al*; Coagulation defects in snake bite poisoning. *J Assoc Physicians India*, 1985;

33 (2): 148-151.

11. Halesha BR *et al*; A study on the clinico-epidemiological profile and the outcome of snakebite victims in a tertiary care center in southern India. *J Clin Diagn Res. Jan* 2013; 7 (1): 122126.
12. Sharma SK *et al*. Impact of snake bites and determinants of fatal outcomes in southeastern Nepal. *Am J Trop Med Hyg.* 2004; 71: 234-238.
13. Saravu K *et al*; Clinical profile, species-specific severity grading, and outcome determinants of snake envenomation: An Indian tertiary care hospital-based prospective study. *Indian J Crit Care Med.*, 2012; 16 (4): 187-192.
14. Inamdar F *et al*; Snakebite: admissions at a tertiary health care center in Maharashtra, India. *S Afr Med J.*, 2010; 100 (7): 456-458.

Incidence and outcome of early post stroke seizures in a tertiary care hospital in New Delhi..... a prospective observational study

Maqsood Ahmad Dar¹, Vasundhara Aggarwal Saika², Joy Dev Mukherjee³, Mukesh Kumar³, Khushbu Goel³

¹ Department of Internal Medicine, GMC Anantnag

² Department of Neurology, PSRI New Delhi

³ Department of Neurology, Max Superspeciality hospital saket New Delhi

ABSTRACT

Background: About 11% of all epilepsies and 30% of newly diagnosed seizures in those older than 60 years have been reported to follow cerebrovascular accidents. Acute symptomatic (early) seizures occur in 3-8% of patients, more commonly in those with severe cortical strokes and intra-parenchymal hemorrhages. This study was undertaken to determine the incidence, risk factors and clinical outcome for early seizures after acute stroke.

Aims and objectives: To study the incidence, risk factors and clinical outcomes of early seizures in post stroke patients in a tertiary care hospital.

Study design: This prospective observational analytical study was conducted from -01-06-2017 to 31-01-2019 in Max Superspeciality Hospital Saket New Delhi. All consecutive patients of acute stroke fulfilling the inclusion and exclusion criteria were included in the study.

Inclusion criteria and exclusion criteria: All patients with acute ischemic or hemorrhagic stroke with age above 18 years, either male or female were included in the study. Patients with a history of seizure or epilepsy before admission, patients with SAH, venous infarct and causes other than vascular origin were excluded from the study.

Results and observation: Out of two hundred and fifty (n=250) patients of acute stroke included in the study early seizures were diagnosed in 29 patients (11.6%).

Conclusions: We concluded that the incidence of post stroke early seizures was 11.6% in our study. We found that Patients with post stroke early seizures had higher NIHSS score, low GCS score and higher modified Rankin score at the time of admission. We recommend that stroke scores like NIHSS, ICH, GCS and mRS should be applied to all the patients of acute stroke to stratify the patients with regard to their risk of developing early seizures and prognosticating the clinical outcome.

Key words: post-stroke, seizures, early, mRS, NIHSS

JMS; 2020;23(3) p?-p? DOI: <https://doi.org/10.33883/jms.v23i3.752>

INTRODUCTION

About 11% of all epilepsies and 30% of newly diagnosed seizures in those older than 60 years have been reported to follow cerebrovascular accidents¹. Acute symptomatic (early) seizures occur in 3-8% of patients, more commonly in those with severe cortical strokes and intra-parenchymal hemorrhages². Late seizures occur in about 8% of patients by 10 years, but most seizures occur in 2 years after an acute Stroke³. This study was undertaken to determine the incidence, risk factors and clinical outcome for early seizures after acute stroke.

Definitions

Stroke is defined according to the WHO's criteria as sudden onset of signs of focal or global disturbance of cerebral function lasting more than 24 hours with no apparent non-vascular cause⁴.

Correspondence:

Dr. Maqsood Ahmad Dar
Neurologist, Department of internal medicine, GMC Anantnag
E mail: drmaq30@gmail.com

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.752>

How to cite this article: Dar MA, Saika V, Mukherjee J, Kumar M, Kumar M, Goel K. Incidence and outcome of early post stroke seizures in a tertiary care hospital in New Delhi..... a prospective observational study. JMS 2020; 23(3):18-23

Received: 2020-05-28 Accepted: 2020-08-29

Early seizures are defined when they appear within the first 7 days after acute stroke onset⁵.

Aims and objectives: To study the incidence, risk factors and clinical outcomes of early seizures in post stroke patients in a tertiary care hospital.

Primary objective: To determine the risk factors of early seizures in post stroke patients.

Secondary objective: To determine the outcome in post stroke patients of early seizures using modified Rankin Scale.

Material and methods

Study population: Consecutive patients of acute stroke admitted in Max Superspeciality Hospital New Delhi, fulfilling the following inclusion and exclusion criteria were included in the study. This prospective observational analytical study was conducted from -01-06-2017 to 31-01-2019.

Inclusion criteria

1. Age above 18 years, either male or female.
2. All the consecutive patients with acute ischemic or hemorrhagic stroke.

Exclusion criteria

1. Patients with a history of seizure or epilepsy before admission.
2. Patients with causes other than vascular origin.
3. Patients with SAH, venous infarct.

Data Collection: Acute stroke patients were subjected to detailed history and clinical examination. Data on vascular risk factors including age, gender, Hypertension (BP>140/90mmhg), Type II Diabetes, dyslipidemias, Ischemic Heart Disease, Atrial Fibrillation, were obtained. All patients underwent CT or MRI brain according to stroke protocol on admission. Stroke severity was evaluated at admission using the National Institute of Health Stroke Scale (NIHSS) and modified Rankin Scale (mRS). Seizures were classified as focal with or without impaired awareness or generalized tonic clonic seizures (GTCS) or non-convulsive seizures (NCSE). Routine investigations including Hemogram, serum biochemistry especially serum Electrolytes, liver function test, kidney function test, random blood sugar, HbA1C to rule out any metabolic abnormality which can be responsible for seizures were

done. Electroencephalogram was done in all patients of stroke with seizures at the earliest possible time.

Methods of measurement of outcome of interest:

Primary outcome: Risk factors for early seizures were identified through logistic regression with seizures and no seizures as dependent variables and risk factors such as Hypertension, Type II Diabetes Mellitus, Hyperlipidemia were taken independent variables. Their odds ratio were worked out along with the 95% confidence interval and statistical significance was at 5% level

Secondary outcome: Clinical outcome were measured including mortality and disability at discharge according to Modified Rankin Score. mRS 0,1,2 was taken as non-disabling stroke and 3,4, as disabling stroke

Statistical Methods: Incidence of early seizures were worked up as percentage of total cases of stroke that will have seizures within 7 days. Risk factors for early seizures were identified through logistic regression with seizures and no seizures as dependent variables and risk factors such as Hypertension, Type II Diabetes Mellitus, Hyperlipidaemia will be independent variables. There odds ratio was worked out along with the 95% confidence interval and statistical significance was at 5% level. Clinical outcomes such as post stroke mortality/ disability was assessed in terms of percentage of cases experiencing early seizures as compared with patients not experiencing seizures. These percentages were compared by Chi-square test. SPSS20 was used for computation.

Results and observation: Total two hundred and fifty (n=250) patients of acute stroke were included in the study. Two hundred twenty patients (88%) had acute ischemic stroke and thirty patients (12%) had acute hemorrhagic stroke. Two hundred and twenty one (n=221) patients (88.4%) of acute stroke did not have early seizures. Early seizures were diagnosed in 29 patients (11.6%). (Figure-1)

Fig. 1: Percentage post stroke early seizures

Table 1: Demographic profile of patients with Acute stroke

Total No. of patients N= 250	Early Seizures		
	Yes (n= 29)	No (n=221)	p Value
Age of patients (Mean $\pm$ SD)	64.83 $\pm$ 16.45	62.13 $\pm$ 14.18	0.396
Sex (M:F)	1.4:1	2.2:1	0.272
Systolic Blood Pressure (mean $\pm$ SD)	146.50 $\pm$ 36.62	143.89 $\pm$ 26.6	0.641
DBP (mean $\pm$ SD)	82.77 $\pm$ 18.49	85.15 $\pm$ 12.3	0.508
Level of Consciousness GCS (mean $\pm$ SD)	8.41 $\pm$ 4.88	14.64 $\pm$ 1.50	< 0.001
CAD	7 (41.7%)	46 (20.8)	0.681
HTN	22 (75.5%)	139 (62.9%)	0.170
AF	4 (7.2%)	16 (13.8%)	0.385
HbA1C (mean $\pm$ SD)	5.94 $\pm$ 1.15	6.27 $\pm$ 1.69	0.328

Risk factors for acute stroke with early seizures:

All the patients were evaluated for preexisting risk factors like glycemic control (HbA1c), presence of coronary artery disease, Type II Diabetes Mellitus, Hypertension, and ECG abnormalities like atrial fibrillation. No significant difference in occurrence of ES was noticed in relation to premorbid risk factors as shown in Table 2.

Table 2: Showing co morbidities of patients included in the study.

HTN	Early seizures				P value
	No		Yes		
	n	%	n	%	
no	82	37.1	7	24.1	0.170
yes	139	62.9	22	75.9	
total	221	100	29	100	
DM					0.573
No	125	56.6	18	62.1	
Yes	96	43.4	11	37.9	
Total	221	100	29	100	
ECG					0.616
Non AF	203	91.8	24	82.7	
AF	18	8.14	5	17.24	
Total	221	100	29	100	
CAD					0.681
No	175	79.2	22	75.9	
Yes	46	20.8	7	24.1	
Total	221	100	29	100	

Ischemic stroke were classified as cortical, subcortical and combined cortical and subcortical based on neuroimaging. Size of the ischemic infarcts was classified as small if lesion is <1.5 cms, medium if lesion was in a cortical superficial branch of MCA or lesion was involving the MCA deep branch, or lesion is in a cortical superficial branch of PCA or lesion was involving the PCA branch or lesion in a cortical superficial branch of ACA. Large if the lesion was involving

the whole territory of ACA or lesion was involving 2 cortical superficial branches of MCA or the lesion was involving a cortical superficial branch of MCA associated to the MCA deep branch or the lesion was involving more than 1 artery territory and lesions involving Brainstem or Cerebellum >1.5 cms. We found that 45.6% had cortical, 29.2% had subcortical and 13.2% had combined cortical and subcortical involvement. When the area of infarct was compared in patients of acute stroke with and without early seizures, it was found that 58.8% of patients of early seizures (n = 10) had combined cortical and subcortical involvement which is statistically significant (p<0.001, Table 3).

Table 3: GCS score of patients with acute stroke with or without seizures at admission

Infarct Regions in MRI/CT	Early Seizure				p value
	No		Yes		
	n	%	N=	%	
Cortical	112	55.2	2	11.8	<0.001
Subcortical	68	33.5	5	29.4	
Cortical+ Subcortical	23	11.3	10	58.8	
Total	203	100	17	100	

We found that 42.2% patients had small infarct, 39.6% had medium size infarct and 12.7% had large infarct on CT /MRI. Seventy percent (70.5%) of patients (n=12) of early seizures had large infarct on CT/MRI which is statistically significant as shown in Figure -2

Fig. 2 :mRS score at admission in patients of acute stroke with and without ES

In patients of hemorrhagic stroke, the size was considered large if the maximal diameter of the lesion was > 3 cms. Out of 12 patients of ICH with ES, 91.6% (n=11) of patients had large bleed which is statistically significant. ICH score was evaluated in patients with hemorrhagic stroke at admission. Out of 30 patients, 12 patients (40%) had ICH score of ≥ 3 . When the comparison was made between the patients with and without seizures, the occurrence of ES was significantly higher (66.66% vs 22.2%) with ICH score of

≥3.(p<0.001.Table-4.

Table 4 : ICH score of patients with acute stroke with and without early seizure

ICH score	Early Seizure				p Value
	No		Yes		
	n	%	n	%	
<3	14	77.7%	4	33.3%	0.001
≥3	4	22.22%	8	66.6%	
Total	18	100%	12	100%	

The most common type of seizure observed was generalized tonic clonic seizure (41.37%). Eleven patients (37.9%) had non-convulsive status epileptics and 6 had focal seizures (20.68%).

GCS score was evaluated in all patients of acute stroke at admission. It was observed that 91.6% had GCS score more than 8 and 8.4% had GCS of ≤ 8. In patients with early seizures, 55.17 % had a score ≤ 8 and 44.8% had score more than 8. The mean GCS score of patients with early seizures was (8.41+ 4.88) and without seizures was (14.64+1.50) which was statistically significant (p<0.001)(Table-5)

Table: 5: GCS score of patients with acute stroke with or without seizures at admission

GCS at admission	Early Seizure				p Value
	No		Yes		
	n	%	n	%	
>8	216	97.7	13	44.8	<0.001
≤8	5	2.26	16	55.17	
Total	221	100	29	100%	

All the patients of acute stroke at admission were evaluated with NIHSS for assessing the severity of stroke. Two hundred and eleven patients (84.4%) had score of less than 16 and 15.6% (n=39) had score of ≥16. When comparison was made between two groups with and without early seizures, it was found that 72.41% of patients with early seizures had a NIHSS score of more than ≥16 at admission and only 8.14% of patients without seizures had the score of ≥16. The mean NIHSS score in patients with early seizures at admission was (13.76 ± 7.68) and those without seizures was (5.77 ± 4.31) which was statistically significant (p<0.001)(Table 6)

Table: 6: NIHSS score in patients of acute stroke with and without seizures

NIHSS at admission	Early Seizure				p Value
	No		Yes		
	n	%	n	%	
<16	203	91.8	8	27.58	<0.001
≥16	18	8.14	21	72.41	
Total	221	100	29	100	

The mRS score was evaluated in all patients of acute stroke at admission. Eighty patients (32%) had mRS score of <2, and 170 patients (68%) had score of ≥2 at admission. When mRS score at admission was compared between patients with early seizures and without seizures, it was found that early seizures occurred in 89.6% of patients of acute stroke with a score of ≥2. These findings were statistically significant (p<0.001- Figure - 3).

Fig. 3 : mRS score at admission in patients of acute stroke with and without ES

Fig. 4: mRS score at discharge of patients of acute stroke

Modified Rankin Score at discharge: Sixty nine patients had disabling stroke (mRS>2) at discharge. Out of these, twenty three patients (79.3%) had ES. The mean mRS at discharge was 41.69 in patients of acute stroke with early seizures as compared to without seizures which is statistically significant (p<0.001) (Table 7).

Table-7: Mean mRS score of patients with post stroke early seizures at discharge

Seizures at discharge							P Value
Early Seizure							
No			Yes				
	Median	Min - Max	Mean ± SD	Median	Min - Max		
mRS at D/S	2.00	1 - 6	4.00 ± 1.69	4.00	1 - 6	<0.001	

Improvement or deterioration of mRS score at discharge :

It was observed that 31% of patients of acute stroke with early seizures had worsening of mRS score at discharge (over a period of 7 days) as compared to only 5% of

patients without seizure group who showed worsening which is statistically significant ($p < 0.001$). Figure -5).

Fig. 5: Improvement and deterioration of mRS of patients with acute stroke with and without ES at discharge.

DISCUSSION:

The rate of early seizures after stroke varies widely in the literature (2.742.8%) due to different inclusion criteria, study designs, and detection modes⁶⁻¹⁴. The Oxford shire community stroke 15 project suggested a cumulative risk of 11.5% for the development of seizures within 5 years. The incidence of early seizures in our study was almost same about 11.6%. However, the reported rate of post stroke early seizures in our study was higher than that reported (2% to 6%) in other prospective studies^{3,2,16}. We did not find any significant association between the presence of co-morbidities like Hypertension, Diabetes mellitus, Ischemic Heart disease, Atrial fibrillation with post stroke early seizures, which is consistent with studies published by Labovitz et al¹⁰ and Lamy et al¹⁶. We found that a higher NIHSS score (>20) increases the risk for the development of seizures. This finding is supported by Bladin et al⁷, Goswami RP et al¹⁷ and Cornado J et al⁷. Szaflarski et al¹⁸ found that the patients with seizures had significantly lower initial Glasgow Coma Scale (GCS) scores, higher National Institutes of Health (NIH) Stroke Scale (NIHSS) score, higher 30-day mortality and had a higher incidence of hemorrhagic stroke than patients without seizure. It was also found that prestroke modified Rankin Score (mRS) >1 was an independent risk factor for development of post stroke seizures. Along with NIHSS score we measured the stroke severity with mRS and GCS at admission. We found that patients with post stroke early seizures had lower GCS score (<10) and higher mRS (≥ 2) at admission in comparison with those patients without seizures. Significant number of patients with seizures had deterioration of mRS in the hospital leading to increase mortality and disability. So in our study low GCS score,

high NIHSS and high Modified Rankin Score were independently associated with development of post stroke seizures.

CONCLUSIONS:

We made following conclusions:

1. The incidence of post stroke early seizures is 11.6% in our study.
2. No association between Age and Gender were found between patients with ES and without ES.
3. We did not find any significant correlation between presence of co morbidities like Hypertension, Diabetes mellitus, Ischemic Heart disease, Atrial fibrillation with occurrence of early seizures in patients of acute stroke.
4. Patients with post stroke early seizures had higher NIHSS score, low GCS score and higher modified Rankin score at the time of admission.

RECOMMENDATION:

We recommend that stroke scores like NIHSS, ICH, GCS and mRS should be applied to all the patients of acute stroke to stratify the patients with regard to their risk of developing early seizures and prognosticating the clinical outcome. This will enable the clinicians to standardize the patient care and treatment including use of prophylactic antiepileptic medications.

Ethical consideration:

This research was carried out in accordance with the basic principles defined in ICMR ' ethical guidelines for biomedical research on human participants (2006); CDSCO' guidance on good clinical practice for clinical research India. The study was conducted only after the approval by the Scientific and Ethics committees of the hospital participating in the study.

REFERENCES:

1. Annegers JF, Hayser WA, O'Brien PC, Whisnant JP. Population based study of seizure disorders after cerebral infarction. *Neurology*. 1996;46:350-355
2. Labovitz DL, Hauser WA, Sacco RL. Prevalence and predictors early seizure and status epilepticus after first stroke. *Neurology*. 2001;57:200-206
3. So EL, Annegers JF, Hauser WA, O'Brien PC, Whisnant JP. Population- based study of seizure disorders after cerebral infarction. *Neurology*. 1996;46:350-355

- 4 (WHO)World health organization Recommendations on stroke prevention, diagnosis and therapy.Report of the WHO Task Force on stroke and other cerebrovascular disorders. Stroke.1989; 20:1407-31
- 5 International League Against Epilepsy Commission on Epidemiology and prognosis, guidelines for epidemiologic studies on epilepsy. Epilepsia 1993;34:592-6
- 6 Burneo, J.G., Fang, J., and Saposnik, G. Impact of seizures on morbidity and mortality after stroke: a Canadian multicenter cohort study. European Journal of Neurology. 2010; 17: 5258
- 7 Bladin CF, Alexandrov AV, Bellavance A et al.Seizures after Stroke: a prospective multicenter study. Arch Neurol. 2000;57:1617-1622
- 8 Berges, S., Moulin, T., Berger, E.,Tatu, L.,Sablot, D., Challier, B. et al. Seizures and epilepsy following strokes: recurrence factors. European Neurology. 2000; 43: 38
- 9 Dodge, P.R., Richardson, E.P., and Victor, M. Recurrent convulsive seizures as a sequel to cerebral infarction: a clinical and pathological study. Brain. 1954; 77: 610646
- 10 Kilpatrick, C.J., Davis, S.M., Tress, B.M.,Rossiter, S.C., Hopper, J.L., and Vandendriesen, M.L. Epileptic seizures in acute stroke. Archives of Neurology. 1990; 47: 157160
- 11 Meyer, J.S., Charney, J.Z., Rivera, V.M., and Mathew, N.T. Cerebral embolization: prospective clinical analysis of 42 cases. Stroke. 1971; 2: 541554
- 12 Lancman, M.E., Golimstok, A., Norscini, J., and Granillo, R. Risk factors for developing seizures after a stroke. Epilepsia. 1993; 34: 141143
- 13 Misirli, H., Özge, A., Somay, G., Erdogan, N., Erkal, H., and Erenoglu, N. Seizure development after stroke. International Journal of Clinical Practice. 2006; 60: 15361541
- 14 Olsen, T.S., Hogenhaven, H., and Thage, O. Epilepsy after stroke. Neurology. 1987; 37: 12091211
- 15 Burn J, Dennis M ,Bamford J ,Sandercock P, Wade D, War low C. Epileptic seizures after a first stroke :The Oxfordshire Community Stroke Project. BMJ. 1997;315: 1582-1587
- 16 Lamy C, Domigo V, semah F. Early and late seizures after cryptogenic stroke in young adults. Neurology.2003;60: 400-404
- 17 Goswami RP1, Karmakar PS,GhoshA.Early seizures in first-ever acute stroke patients in India: incidence, predictive factors and impact on early outcome.Eur JNeurol. 2012 Oct; 19(10): 1361-6.
- 18 Szaflarski, J.P., Rackley, A.Y. Kleindorfer, D.O., Khoury, J., Woo, D., Miller, R. et al. Incidence of seizures in the acute phase of stroke: a population based study. Epilepsia. 2008; 49: 974981.

ORIGINAL ARTICLE

Effect of Metformin on TSH Levels

Idrees Mubarak¹, Shariq Rashid Masoodi², Zafar Amin Shah³, Syed Mudassar⁴,
Saba Moosaraza⁵, Mir Fawad Ahmed⁶

¹Department of Endocrinology, Aster Hospital Dubai, ²Department of Endocrinology, SKIMS

³Department of immunology, SKIMS, ⁴Department of Biochemistry, SKIMS

⁵Department of Anesthesiology, Sita Bhateja Hospital Bangalore, ⁶GMC Srinagar

ABSTRACT

Aims and objectives: Main aim of the study was to see the effect of metformin on thyroid function parameters especially TSH. We also wanted to see factors responsible for these thyroidal effects of metformin.

Materials and methods: we selected treatment naïve type 2 diabetes and PCOS subjects. we had 72 subjects, 56 were having type 2 diabetes and 16 were having PCOS. We took their blood samples at baseline before starting metformin and 6 months after starting metformin. In both samples we checked thyroid functions and insulin resistance indices. Dose of metformin was 1 gm in type 2 DM and 500mg PCOS. We took 80 patients at beginning, 8 were lost to follow up, 72 completed the study.

Results: we found that there was no significant change in thyroid function parameters after 6 months of metformin intake. It made no difference whether patients were obese or non-obese, diabetic or nondiabetic, male or female. But on subgroup analysis it was found that in those patients in whom insulin resistance decreased, TSH was significantly decreased where as in those subjects in which insulin resistance had increased TSH did not change significantly.

Conclusion: In this study we concluded that metformin decreases TSH especially in those patients where it decreases insulin resistance also successfully.

Clinical Significance: Minor elevations in TSH may be part of metabolic syndrome and may not warrant treatment with thyroxine.

Key words: Metformin, TSH

JMS: 2020; 23(3): DOI: <https://doi.org/10.33883/jms.v23i2.765>

INTRODUCTION

Type 2 diabetes mellitus (DM) is the predominant form of diabetes mellitus worldwide, accounting for 90% of cases globally.^{1,2} Insulin resistance is a consistent finding in patients with type 2 DM, which may be present for years before the onset of diabetes.^{3,7} Prospective studies show that insulin resistance predicts the onset of diabetes mellitus.^{3,4} Polycystic ovary syndrome (PCOS), a heterogeneous disorder of unclear etiology, when fully expressed, the manifestations include ovulatory dysfunction, androgen

excess, polycystic ovaries, and obesity.⁸ Burghen et al, 1980 were among the first to report that PCOS patients were hyperinsulinemic in response to an oral glucose tolerance test.⁹ It is now known that insulin resistance, and the development of compensatory hyperinsulinemia, is a

Correspondence:

Dr. Idrees Mubarik, D.M (Endocrinology)
Specialist Endocrinologist
Aster Hospital Dubai
Phone : +971565623704
Email: mubarikidrees@gmail.com

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.773>

How to cite this article:

Received: 2020-05-25 Accepted: 2020-06-12

frequent finding in PCOS.¹⁰

Metformin remains a time tested drug and stands as a drug of first choice in majority of the patients with type 2 DM. It is also frequently used as a treatment of PCOS especially in obese and glucose intolerant patients. In addition to antihyperglycemic action metformin has a pleiotropic effect which is exerted by metformin on arterial vessels by its effects on lipids, inflammation, hemostasis, endothelial and platelet function and vessel wall abnormalities.¹¹

One of the latest domains of function added to metformin is on thyroid functions especially TSH. Few years back Robert et al, 2006 reported that metformin appeared to suppress serum TSH to subnormal levels without resulting in biochemical or clinical hyperthyroidism.¹² Rotondi et al, also showed that metformin results in thyrotropin suppression in patients of PCOS.¹³

In this study we tried to see the effect of metformin on thyroid functions in patients of type 2 diabetes mellitus and PCOS who had never received this drug before.

PATIENTS AND METHODS

This study was conducted in the Department of Endocrinology at Sher-i-Kashmir Institute of Medical Sciences (SKIMS), Srinagar. Study was approved by the Institute Ethical Committee. Patients were recruited prospectively over a period of 1 year; each patient was part of the study for a period of 6 months.

Subjects included (1) newly diagnosed type 2 diabetes mellitus patients as per ADA statement 2013¹⁴, who were treatment naïve and (2) first time diagnosed patients of PCOS as per Rotterdam criteria 2003.¹⁵

Patients with overt hyperthyroidism or hypothyroidism were excluded from study and so were the patients who had taken metformin earlier for one reason or the other.

All the patients were put on metformin with a dose of 1gm once daily in type 2 diabetes mellitus and 500 mg once daily in PCOS patients. Height, weight, BMI and abdominal circumference of all subjects were measured at the recruitment. We measured total T3, total T4, free T3, free T4, TSH, anti-TPO antibody, serum insulin, blood sugar fasting levels at baseline and after 6 months of treatment with metformin. Lipid profile, serum creatinine, and uric acid were also measured at baseline. Insulin was measured

by immunoradiometric assay (IRMA). Analytical sensitivity of assay was 0.49µIU/ml. The range of measurement was from 0.49-300 µIU/ml. Anti-TPO antibody was measured by ELISA method. Values less than 50 IU/ml were considered negative, values between 50-75 IU/ml were considered intermediate and values more than 75 IU/ml were considered positive.

TSH was measured by hypersensitive third generation chemiluminescent immunoassay, with a range of detection from 0.01-100 µIU/ml. coefficient of variation ≤ 20%. Normal range in this assay is 0.34-5.6 uIU/ml.

Free T3, Free T4, total T3 and total T4 were measured by chemiluminescent immunoassay on Beckman Coulter immunoassay system. Patient test results are determined automatically by the system software using a weighted four parameter logistic curve (4PLC) math model. The range of detection for free T3 was 0.88 to 30 pg/ml. This assay exhibits total imprecision ≤ 12%. Normal range of free T3 in this assay is established as 2.5-3.9 pg/ml. Range of detection for free T4 was 0.256-0 ng/dL. Normal range for free T4 with this assay is established as 0.61-1.12 ng/dl. For total T3 range of detection was 0.1-8 ng/ml, with imprecision less than 10%. Established normal range for total T3 is 0.87-1.78 ng/ml. Range of total T4 detection was 0.5-30 µg/dl with an imprecision of less than 10%. Established normal range for total T4 by this assay is 6.09-12.23 µg/dl.

We also calculated HOMA-IR and QUICKI indices using standard formulas. $HOMA-IR = \text{Fasting insulin (mIU/ML)} \times \text{Fasting glucose (mg/dl)} / 405$,

$QUICKI = 1 / (\log \text{fasting insulin} + \log \text{fasting glucose})$

RESULTS

Of the 80 subjects recruited, only 72 completed the study, but the characteristics of 8 subjects who didn't follow were similar to those who completed the study. Of these 72 study subjects, 56 were patients of type 2 DM while the remaining 16 subjects were having PCOS. All the subjects were diagnosed first time and were treatment naïve. Age of the subjects ranged from 17 to 65 with a mean of 40.76 ± 11.72 years (median, 43). Understandably subjects with PCOS were significantly younger than those with type 2 DM. Relevant parameters at baseline are shown in table 1.

Table 1: Baseline anthropometric and laboratory parameters of the study subjects

Parameter	T2DM (mean ± SD)	PCOS (mean ± SD)	All subjects
Age (years)	46.79 ± 2.21	23.19 ± 3.66	40.76 ± 11.72
BMI (Kg/m ²)	25.95 ± 2.35	22.69 ± 2.06	25.23 ± 2.65
HOMA IR	4.91 ± 4.02	2.86 ± 2.04	4.45 ± 3.76
QUICKI	0.32 ± 0.04	0.35 ± 0.06	0.33 ± 0.05
Cholesterol (mg/dl)	193.86 ± 25.53	173.56 ± 23.40	189.35 ± 26.32
Triglycerides (mg/dl)	215.14 ± 37.13	175.19 ± 28.04	206.26 ± 38.90
LDL (mg/dl)	112.64 ± 18.45	98.81 ± 12.31	109.57 ± 18.15
HDL (mg/dl)	43.64 ± 4.047	46.25 ± 3.73	44.22 ± 4.10
T3 (ng/ml)	1.07 ± 0.26	1.19 ± 0.22	1.10 ± 0.25
T4 (µg/dl)	8.59 ± 1.80	7.40 ± 0.88	8.32 ± 1.71
TSH (uIU/ml)	3.38 ± 2.10	3.10 ± 1.54	3.32 ± 1.98
FT3 (pg/ml)	2.78 ± 0.31	3.05 ± 0.29	2.84 ± 0.32
FT4 (ng/dl)	0.91 ± 0.16	0.79 ± 0.08	0.88 ± 0.16

We observed that TSH was having a positive linear relation with HOMA IR, more the insulin resistance, higher were the TSH levels.

FIGURE 1: Positive linear relationship between TSH and HOMA IR ($r=0.4$, $p=0.03$)

After 6 months of treatment with metformin the mean TSH values were 2.9 ± 1.64 μ IU/ml. Mean total T4 levels were 7.54 ± 1.26 μ g/dl and mean total T3 levels were 1.12 ± 0.26 ng/ml. Mean free T4 levels were 0.84 ± 0.11 ng/dl, while mean free T3 levels were 2.83 ± 0.31 pg/ml (figure 2). Mean HOMA-IR was 3.41 ± 2.66 and mean QUICKI was 0.198 ± 0.015 .

Overall the mean TSH values decreased from 3.32 to 2.9 μ IU/ml after treatment with metformin, but the difference was not statistically significant when compared by paired t test ($p=0.07$). However, total T4 values decreased

significantly from 8.32 to 7.5 μ g/dl ($p=0.03$). There was no significant difference between baseline total T3 and post therapy total T3 ($p=0.548$). There was also no difference between basal and post therapy free T3 ($p=0.848$). Again there was no difference between basal and post therapy free T4 levels ($p=0.128$). (table 2)

Table 2: Thyroid function parameters and insulin resistance markers before and after treatment.

Parameter	Baseline	After 6 months	p value
Total T3 (ng/ml)	1.10 ± 0.25	1.12 ± 0.26	0.55
Total T4 (µg/dl)	8.32 ± 1.71	7.55 ± 1.26	0.03
Free T4 (ng/dl)	0.88 ± 0.16	0.85 ± 0.11	0.13
Free T3 (pg/ml)	2.84 ± 0.32	2.83 ± 0.32	0.85
TSH (µIU/ml)	3.32 ± 1.98	2.90 ± 1.65	0.07
QUICKI	0.33 ± 0.05	0.33 ± 0.04	0.18
HOMA-IR	4.45 ± 3.76	3.41 ± 2.66	<0.01

Figure 2 A comparison of change in TSH between two groups where HOMA-IR decreased and increased respectively.

DISCUSSION

The primary aim of present study was to observe effect of metformin on TSH. We also tried to assess the relation between indices of insulin resistance and parameters of thyroid function, especially TSH. We also tried to find whether any thyroidal effects of metformin were related to changes in markers and/or indices of insulin resistance.

We selected the first time diagnosed patients of type 2 DM and PCOS who were historically euthyroid and not taking any medications that could affect TSH levels. Age of subjects ranged from 17 to 65 years. Mean age for T2DM patients was 45.79 ± 7.63 years whereas that of PCOS subjects was 23.19 ± 3.66 years. Out of 72 subjects (40 females), 39 (54%) had BMI of 25 kg/m^2 or more. Baseline HOMA-IR of our subjects ranged from 0.2 to 16.2, with a mean of 4.45 ± 3.76 (median, 3.4), with a significant linear correlation with age ($r = 0.33$) and body weight ($r = 0.24$). Baseline QUICKI ranged from 0.26 to 0.51 with a mean of 0.33 ± 0.05 (median, 0.32); correlating significantly with age ($r = -0.29$; $p = 0.014$) only. Correlation of body weight and BMI with HOMA-IR is a well-known fact proven beyond doubt in many studies.¹⁶

The mean TSH of our study subjects at baseline was $3.32 \pm 1.98 \text{ } \mu\text{IU/ml}$, which corresponds well to the range for normal euthyroid population.¹⁷ There was a positive linear correlation between TSH and markers of insulin resistance (HOMA-IR) especially in those subjects who had free T4 in upper half of normal range. Similar correlations have been shown in previous studies. In a recent study from South-India, Vyakaranam et al, reported that TSH positively correlated with insulin and HOMA-IR.¹⁸ Singh et al, also have noticed this positive correlation of TSH with indices of insulin resistance, but only in overt hypothyroid group of their study and not in euthyroid and subclinical hypothyroidism subjects.¹⁹ Their study was primarily aimed at observing this correlation only. In another study association of TSH with insulin resistance and androgen parameters in PCOS patients was studied by Dittrich R et al.²⁰ The authors observed that women with TSH of $>2.5 \text{ mU/l}$ had higher BMI, higher insulin and higher testosterone concentrations. In another study higher levels of TSH were found to be associated with higher LDL cholesterol and lower HDL cholesterol levels and thereby

were associated with increased cardiovascular risk.²¹

After 6 months of therapy with metformin we found an overall decrease in mean TSH levels (3.32 vs $2.91 \text{ } \mu\text{IU/ml}$). The level of suppression we observed was just marginally short of statistical significance ($p=0.07$). There are studies though, that have observed a statistically significant suppression of TSH by metformin. Carlo Capelli et al, in their study observed that metformin causes suppression of TSH in hypothyroid patients, both treated and untreated, but they failed to show this effect in euthyroid subjects.²² They hypothesized that metformin may increase the inhibitory modulation of T4 on TSH secretion and this effect may not be valid in case of a normal hypothalamo-pituitary-thyroid axis, thus explaining the lack of effect on euthyroid patients where axis is normally operating. Mario Rotondi et al, in their study on PCOS subjects with and without hypothyroidism observed that metformin caused TSH suppression of statistical significance in only hypothyroid subjects and had no effect on TSH in euthyroid subjects.²³ Vigersky et al, reported suppression of TSH to below normal in 4 hypothyroid patients who were on treatment and well controlled, but there were no clinical features of thyrotoxicosis in these patients.²⁴ Another reason that in our study suppression of TSH did not achieve statistical significance may have been a relatively small sample size.

One unique observation from our study is significant decrease in total T4 after 6 months of treatment with metformin, but there was no change in free T4. None of the studies mentioned above have noticed a suppression in total T4. This effect noticed in our study may be due to possible effect of metformin on the protein binding of thyroxine as has been postulated before also.^{24,25}

We did not find any difference in change in thyroid function on the basis of their BMI (obese vs non-obese), gender (male vs female) and diagnosis (T2DM vs PCOS). Oleadri SE et al, in their study on subjects with abdominal obesity did not find any variation in TSH levels on treatment with metformin. There is no study in literature that has compared head to head, changes in TSH among males vs females, or diabetes vs PCOS.

We also observed that TSH was significantly decreased in those subjects in whom HOMA-IR has decreased as compared to those subjects where HOMA-IR has increased.

Though there are no such studies till date who have shown correlation of thyroidal effect of metformin with changes in HOMA-IR, but many studies have shown TSH to be related to insulin resistance.¹⁸⁻²¹ These studies have shown more insulin resistance in subclinical hypothyroidism as compared to euthyroid population.¹⁸ Singh et al, found that TSH progressively increased through low to high BMI and through increasing levels of serum insulin.¹⁹ Ours is the first study which shows that TSH suppression of metformin may be dependent on decrease in insulin resistance.

There have been various hypotheses regarding the mechanism of action of metformin in the context of TSH suppression, which include:²⁵⁻²⁶ 1) Metformin may increase the affinity or number of thyroid hormone receptors 2) There may be an increase in dopaminergic tone resulting in TSH suppression 3) Change in bioavailability of thyroid hormone by altering protein binding 4) Metformin may cause constitutive activation of TSH receptor. However, to elucidate the exact mechanism of action is beyond the scope of current study.

The positive points of our study are that it is a prospective study as compared to majority of the studies mentioned previously, which were retrospective in nature. Also in our study, participants were not taking any other medication except metformin, where as in other studies patients were on multiple other drugs including thyroxine. Majority of our patients tested negative for anti-TPO antibody, only 5 out of 72 tested positive. This makes possibility of autoimmune influence on the thyroid functions less likely; any change in TSH would be more likely due to time and/or metformin. Majority of previous studies have not checked for anti-TPO antibody. We have used baseline values as control though, it would have been ideal to have separate controls. Another limitation of our study is a marginally small sample size.

CONCLUSION

In conclusion, we observed a positive correlation between serum thyrotropin levels and indices of insulin resistance. There was a decrease of thyrotropin (TSH) level with metformin treatment at 6-months follow up that is statistically significant only in subjects in whom HOMA-IR also decreased. We also found that total T4 levels were decreased significantly after treatment with metformin, but this effect was not dependent on change in HOMA-IR, and needs to be studied further. How exactly the changes in

HOMA-IR bring about changes in TSH levels remains a topic of interest for further research.

REFERENCES

1. Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic. *Nature* 2001; 414:782-787.
2. King H, Aubert RE, Herman WH. Global burden of diabetes, 1995-2025: prevalence, numerical estimates, and projections. *Diabetes Care* 1998; 21:1414-1431.
3. Warram JH, Martin BC, Krowelski AS, Soeldner JS, Kahn CR. Slow glucose removal rate and hyperinsulinemia precede the development of type II diabetes in the offspring of diabetic parents. *Ann Intern Med* 1990; 113:909-915.
4. Lillioja S, Mott DM, Howard BV, Bennett PH, Yki-Järvinen H, Freymond D, Nyomba BL, Zurlo F, Swinburn B, Bogardus C. Impaired glucose tolerance as a disorder of insulin action: longitudinal and cross-sectional studies in Pima Indians. *N Engl J Med* 1988; 318:1217-1225.
5. Haffner SM, Stern MP, Dunn J, Mobley M, Blackwell J, Bergman RN. Diminished insulin sensitivity and increased insulin response in nonobese, nondiabetic Mexican Americans. *Metabolism* 1990; 39:842-847.
6. Reaven GM, Bernstein R, Davis B, Olefsky JM. Non-ketotic diabetes mellitus: insulin deficiency or insulin resistance? *Am J Med* 1976; 60:80-88.
7. DeFronzo RA. Lilly lecture 1987. The triumvirate: beta-cell, muscle, liver. A collusion responsible for NIDDM. *Diabetes* 1988; 37:667-687.
8. Stein IF, Leventhal NL. Amenorrhea associated with bilateral polycystic ovaries. *Am J Obstet Gynecol* 1935; 29:181-185
9. Burghen GA, Givens JR, Kitabchi AE. Correlation of hyperandrogenism with hyperinsulinism in polycystic ovarian disease. *J Clin Endocrinol Metab* 1980; 50:113-116
10. DeUgarte CM, Bartolucci AA, Azziz R. Prevalence of insulin resistance in the polycystic ovary syndrome using the homeostasis model assessment. *Fertil Steril* 2005; 83:1454-1460
11. Anfossi G, Russo I, Bonomo K, Trovati M., The

- cardiovascular effects of metformin: further reasons to consider an old drug as a cornerstone in the therapy of type 2 diabetes mellitus. *Curr Vasc Pharmacol*. 2010; 8(3):327-337.
12. Vigersky RA, Filmore-Nassar A, Glass AR. Thyrotropin suppression by metformin, a brief report. *J Clin Endocrinol Metab*, 2006;91(1):225-227.
13. Rotondi M, Cappelli C, Magri F, Botta R, Dionisio R, Iacobello C, De Cata P, Nappi RE, Castellano M, Chiovato L. Thyroidal Effect of Metformin Treatment in Patients With Polycystic Ovary Syndrome. *Clin Endocrinol* 2011;75(3):378-381.
14. American Diabetes Association. Executive summary: standards of medical care in diabetes 2013. *Diabetes Care* 36: S4-S10.
15. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group (2004) Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril*. 2004;81(1):19-25.
16. Panag KM, Kaur N, Goyal G. Correlation of insulin resistance by various methods with fasting insulin in obese. *Int J Appl Basic Med Res* 2014;4(Suppl 1):S41-S45.
17. Spencer CA, Schwarzbein D, Guttler RB, LoPresti JS, Nicoloff JT. Thyrotropin (TSH)-releasing hormone stimulation test responses employing third and fourth generation TSH assays. *J Clin Endocrinol Metab*. 1993;76(2):494-8
18. Vyakaranam S, Vanaparthi S, Nori S, Palarapu S, Bhongir AV. Study of Insulin Resistance in Subclinical Hypothyroidism. *Int J Health Sci Res* 2014; 4(9): 1471-153.
19. Singh BM, Goswami B, Mallika V. Association between insulin resistance and hypothyroidism in females attending a tertiary care hospital. *Indian J Clin Biochem* 2010;25(2):141-5.
20. Dittrich R, Kajaia N, Cupisti S, Hoffmann I, Beckmann MW, Mueller A. Association of thyroid-stimulating hormone with insulin resistance and androgen parameters in women with PCOS. *Reprod Biomed Online*. 2009;19(3):319-25.
21. Bakker SJ, ter Maaten JC, Popp-Snijders C, Slaets JP, Heine RJ, Gans RO. The relationship between thyrotropin and low density lipoprotein cholesterol is modified by insulin sensitivity in healthy euthyroid subjects. *J Clin Endocrinol Metab*. 2001;86(3):1206-11.
22. Cappelli C, Rotondi M, Pirola I, Agosti B, Gandossi E, Valentini U, De Martino E, Cimino A, Chiovato L, Agabiti-Rosei E, Castellano M. TSH-Lowering Effect of Metformin in Type 2 Diabetic Patients Differences between euthyroid, untreated hypothyroid, and euthyroid on L-T4 therapy patients. *Diabetes Care* 2009;32(9):1589-90.
23. Rotondi M, Cappelli C, Magri F, Botta R, Dionisio R, Iacobello C, De Cata P, Nappi RE, Castellano M, Chiovato L. Thyroidal effect of metformin treatment in patients with polycystic ovary syndrome. *Clinical Endocrinology* 2011;75(3):378-81
24. Vigersky RA, Filmore-Nassar A, Glass AR. Thyrotropin Suppression by Metformin. *J.Clin.Endocrinol.Metab* 2006;91(1):225-7.
25. Theodora Pappa, Maria Alevizaki. Metformin and Thyroid: An Update. *Eur Thyroid J* 2013;2:222-8
26. Chau-Van C, Gamba M, Salvi R, Gaillard RC, Pralong FP. Metformin inhibits adenosine 5'-monophosphate-activated kinase activation and prevents increases in neuropeptide Y expression in cultured hypothalamic neurons. *Endocrinology*. 2007;148(2):507-11.

Recurrent Small Gut Fecalomas Causing Obstruction- A Case Report

Mudassir Ahmad Khan ¹, Asif Mehraj ², Arshad Baba ³, Mushtaq Laway ³, Rauf A Wani ², Fazl Q Parry ², Nisar A Chowdri ²

¹ Department of General Surgery, GMC Rajouri

² Department of General Surgery, Colorectal Surgery Division, SKIMS

³ Directorate of Health Services Kashmir

ABSTRACT

Fecaloma is a mass of hardened faeces difficult to pass through the intestinal lumen. Due to various reasons, a bolus of fecal matter stagnates, accumulates and inspissates in the intestinal lumen, until it becomes big enough to cause intestinal obstruction. Though there are various predisposing factors for the formation of fecaloma but the actual cause of fecaloma formation is still unknown. Small gut fecalomas are very rare. Herein we are presenting a case of 50 year male who underwent laparotomy three times for small gut obstruction caused due to recurrent formation of small gut fecaloma.

Keywords: Fecaloma; Small gut obstruction; Enterotomy; Laparotomy; Recurrent.

JMS: 2020; 23(3): DOI: <https://doi.org/10.33883/jms.v23i3.763>

BACKGROUND

First described by Abella ME in 1967¹, fecaloma is a mass of hardened faeces difficult to pass through the intestinal lumen. Due to various reasons a bolus of fecal matter may stagnate and accumulate in the intestinal lumen, then it may get inspissated and gradually increase in size until the intestine may become obstructed. Fecaloma physically appears like a tumor²⁻³ on palpation of the gut intra-operatively. Fecaloma formation has been associated to various diseases like Hirschsprung's disease⁴, psychiatric patients, Chagas disease, spinal cord injuries, secondary to inflammatory and neoplastic diseases, and in chronic constipation⁵. Most of the impacted fecalomas have been reported in rectum and sigmoid⁶⁻⁷. Up till now only few cases of small gut fecaloma have been reported in the literature⁸⁻⁹. Herein we have a case to report of a patient who got operated many times for recurrent small gut fecaloma obstruction. To our knowledge it is the first case report of recurrent small gut fecaloma obstruction.

Case Description: This is an interesting case of a 50-year

man with a surgically significant past history. About 20 years back the patient had undergone laparotomy with gastrojejunostomy with vagotomy for chronic duodenal ulcer and an emergency open appendectomy was done about 8 years ago. The patient was referred to our emergency room with abdominal pain, vomiting and abdominal distention for 3 days. Diagnosis of Intestinal obstruction was made and E/L with enterotomy with removal of a large fecaloma (4x8 cm large jejunal fecaloma; mixed with dead worms) at about 50cm from DJ. Also large gut contained multiple small (1x1cm) fecalomas. Post-op period was uneventful and patient was discharged on 7th POD.

Almost after one year, patient again got operated for small gut obstruction (Fig. 1) with E/L with adhesionolysis with

Correspondence:

Dr. Mudassir Ahmad Khan, MS
Assistant Professor, Department of General Surgery,
GMC Rajouri
Email: khanmudassir925@gmail.com

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.763>

How to cite this article: Khan MA, Mehraj A, Baba A, Laway M, Wani RA, Parry FQ, Chowdhary NA. Recurrent Small Gut Fecalomas Causing Obstruction- A Case Report. JMS 2020; 23(3):30-32

Received: 2020-06-28 Accepted: 2020-08-11

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

enterotomy with removal of fecaloma. This time intra-operatively there were multiple dense adhesions between gut loops with moderate amount of inter-loop fluid, hard fecaloma impacted at about 3 feet proximal to ileo-Caecal valve, gut proximal to fecaloma dilated and edematous wall. Post-op period un-eventful and patient got discharged on 7th POD. After about two years of this second laparotomy for obstruction, patient again developed the features of small bowel obstruction which was initially managed conservatively as a case of SAIO but in view of recurrent episodes of SAIO the patient was ultimately managed by diagnostic lap followed by exploratory laparotomy with adhesiolysis with enterotomy with evacuation of multiple fecalomas with loop ileostomy. Intra-operatively dense adhesions involving small bowel were present with multiple fecalomas present in small bowel with dilatation of small gut proximal to the fecaloma impaction. Due to very dense adhesions the ileostomy loop could not be mobilized properly and we landed up in creation of proximal loop ileostomy that too not at a proper location and instead needed to be confectioned in the midline at supraumbilical region in the main laparotomy wound (Fig. 2). Post-operatively the patient managed conservatively for fever with antibiotics and was discharged on 9th POD.

Fig. 1: CT picture of the patient showing features of small bowel obstruction.

Fig. 2: Ileostomy in the midline at supraumbilical region

Because of history of recurrent small gut fecaloma-obstruction and before taking down the ileostomy we tried to find out any systemic cause for the formation of recurrent fecalomas in this patient. We did his thyroid profile, CFTR gene mutation (for adult onset cystic fibrosis) and CT enterography. However, all these tests were normal. Colonoscopy was done up to hepatic flexure that showed features of diversion colitis. Enteroscopy done via the stoma showed that the proximal jejunal loop was normal, distal ileal loop examined only up to 30cm which was also normal (because the scope could not be advanced due to loop formation), conray injected through scope did not reveal any obstruction. Ultimately after about 6 months of last operation, the ileostomy was taken down unevent-fully and patient was put on low fibre diet with laxatives and till now patient is doing well.

DISCUSSION

Since only few cases of fecaloma causing small gut obstruction have been reported in literature, the exact incidence and prevalence of occurrence of small gut fecalomas is not known. Henceforth the treatment options are not standardized and vary with the factors like level of obstruction, condition of obstructed bowel and presence of co-morbidities. Stool softeners may be used by rectal route and even the digital rectal evacuation for recto-sigmoid fecalomas and oral mineral and olive oil usage are appropriate treatment methods for mild and moderate obstructions due to fecalomas¹⁰. However, when medical treatment is unsuccessful or when multiple impassable fecalomas are present or when the patient presents with features of acute obstruction, the surgical intervention becomes must. The recto-sigmoid is the most commonly involved area by fecalomas and the small bowel is a most unusual site¹¹. This may be explained by the fact that stools in the left colon are firmer and have more solid consistency than those in other regions and also because the colon diameter is narrower on the left than the right^{2&12}.

Coming to the discussion of our present case, the fecalomas developed recurrently in the small intestine landing the patient three times into emergency theatre for laparotomy and enterotomy and finally also the covering ileostomy. His thyroid profile, CFTR gene mutation, CT enterography, enteroscopy and colonoscopy were normal and apparently we could not find any cause for the formation of recurrent fecalomas in this patient. However the patient had a dietary

history of consuming excessive maize in his diet. Does any particular diet predisposes to fecaloma formation in the small intestine, is to be studied. Also the patient was a follow up case of laparotomy for duodenal ulcer treatment. Besides causing SAIO, does the post-operative adhesions predispose to the formation of small gut fecalomas is another query to be answered. The reason for 'trouble-causing recurrent small gut fecaloma formation' in our patient still remains unanswered. However till now, the patient is doing well but God knows better when he will have adhesion or fecaloma obstruction again.

CONCLUSION

Only few cases of fecaloma causing small gut obstruction have been reported in literature, the exact incidence and prevalence of occurrence of small gut fecalomas is not known. Fecaloma formation may land the patient in trouble of getting operated in emergency settings for intestinal obstruction. The cause of recurrent formation of small gut fecalomas remains unknown. Our case needed to be operated three times for recurrent fecaloma obstruction and fourth time for ileostomy take-down.

Clinical Significance: Fecaloma formation may land the patient in trouble of getting operated in emergency for intestinal obstruction. The actual cause for recurrent formation of small gut fecalomas remained unknown.

Acknowledgment: For managing this case very efficiently, we acknowledge the support rendered by the resident doctors and the nursing staff especially the Stoma Care Nurses of the Colorectal Division of the Department of General Surgery SKIMS, Soura Srinagar.

Conflict of Interest: Authors declare that they have no conflict of interest.

Funding declaration: No funding was received in preparation of this manuscript.

REFERENCES

- 1) Abella ME, Fernández AT. Large fecalomas. *Dis Colon Rectum*. 1967;10(5):401-4.
- 2) Yucel AF, Akdogan RA, Gucer H. A giant abdominal mass: fecaloma. *Clin Gastroenterol Hepatol*. 2012;10(2):e9-e10.
- 3) Zurabishvili K, Rekhviashvili A, Sakhamberidze M, Tsiklauri K. A case of giant fecaloma in a 24-yearold woman. *Georgian Med News*. 2015(240):11-4.
- 4) Campbell JB, Robinson AE. Hirschsprung's disease presenting as calcified fecaloma. *Pediatr Radiol*. 1973;1(3):161-3.
- 5) Caiazzo P, De Martino C, Del Vecchio G, Di Lascio P, Marasco M, Laviani F, et al. Megacolon for a giant faecaloma with unlucky outcome: case report and review of the literature. *Ann Ital Chir*. 2013;84(3):319-22.
- 6) Rajagopal A, Martin J. Giant fecaloma with idiopathic sigmoid megacolon: report of a case and review of the literature. *Dis Colon Rectum*. 2002;45(6):833-5..
- 7) Zimmers T. Giant calcified fecaloma. *Ann Emerg Med*. 1984;13(9 Pt 1):749.
- 8) Mushtaq M, Shah MA, Malik AA, Wani KA, Thakur N, Parray FQ. Giant Fecaloma Causing Small Bowel Obstruction: Case Report and Review of the Literature. *Bull Emerg Trauma*. 2015;3(2):70-72.doi: 10.7508/beat.2015.02.007.
- 9) Ha Yeong Yoo, Hye Won Park, Seong-Hwan Chang and Sun Hwan Bae. Ileal Fecaloma Presenting with Small Bowel Obstruction. *Pediatr Gastroenterol Hepatol Nutr* 2015 September 18(3):193-196. pISSN: 2234-8646 e I S S N : 2234-8840 <http://dx.doi.org/10.5223/pghn.2015.18.3.193>
- 10) Aykut Soyder, Hedef Özgün. Giant Fecaloma in a 16-Year-Old Boy: Case Report. *Kolon Rektum Hast Derg* 2012;22:160-163
- 11) Cid AA, Pietruk T, Bidari CZ, Ehrinpreis MN. Cecal fecaloma mimicking colonic neoplasm. *Dig Dis Sci*. 1981;26:1134-7.
- 12) Sakai E, Inokuchi Y, Inamori M, Uchiyama T, Iida H, Takahashi H, et al. Rectal fecaloma: successful treatment using endoscopic removal. *Digestion* 2007;75:198.

CASE REPORT

Case of Van-Wyk Grumbach syndrome

Vishnu Vasudevan, Arun Viswanath, Riaz Ahmad Misgar

Department of Endocrinology, Sher I Kashmir Institute of Medical Sciences, Srinagar, India

ABSTRACT

Background: Hypothyroidism in childhood typically leads to delayed puberty, rarely can it cause precocious puberty - this latter complex is eponymously known as Van Wyk-Grumbach syndrome.

Case description: We report a case of a seven and a half year old girl who was referred with vaginal bleeding after after starting her on levothyroxine replacement therapy for newly detected primary hypothyroidism. This case report illustrates the typical features of this rare syndrome for primarily educational purposes. Many case reports of Van Wyk-Grumbach syndrome have been published in world literature over the past seven decades after the initial description. Though the features of this syndrome are fairly well described in the literature, the understanding of pathophysiological mechanisms is speculative and still evolving.

Clinical relevance: Wide spread awareness of the syndrome will encourage primary physicians and paediatricians to counsel the family of a child before starting treatment for hypothyroidism which will help avoid unnecessary worry and more importantly inadvertent procedures or intervention.

JMS: 2020; 23(3): DOI: <https://doi.org/10.33883/jms.v23i3.777>

INTRODUCTION

Van Wyk and Grumbach first described this eponymous syndrome 60 years back as an association of hypothyroidism with precocious puberty (PP) and ovarian mass.¹ They mentioned it as an example of hormonal overlap in pituitary feedback; which is still an accepted pathophysiological mechanism for this peculiar syndrome. It causes gonadotropin independent iso-sexual precocious puberty and manifests as such in female children; males may be affected too causing macroorchidism. Thyroid hormone replacement causes reversal of almost all of the manifestations and it is crucial to consider this possibility in a girl child presenting with breast enlargement, adnexal mass or menstrual bleeding at an earlier age than normal.²

Here we describe a girl child who was referred to our endocrinology out-patient department with menstrual bleeding after starting treatment for hypothyroidism.

CASE REPORT

A 7 years and 4 months old girl was referred to our endocrine OPD with vaginal bleed since 4 days. She was from a poor socioeconomic background, the product of a non-consanguineous marriage, third in birth order and was born at term after an uneventful antenatal period. Her birth weight is unknown but as a new-born she was hospitalized for 3 days for some feeding difficulties (details unavailable). There is no history of prolonged jaundice, hypoglycaemia, seizures, umbilical hernia, constipation or hypotonia. She was breastfed till 2 years and achieved all milestones at appropriate age. She was apparently normal

Correspondence:

Dr Raiz Ahmad Misgar,
Associate Professor, Department of Endocrinology, SKIMS
Email: drreyaz07@rediffmail.com

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.777>

How to cite this article: Vasudevan V, Viswanath A, Misgar RA. Case of Van-Wyk Grumbach syndrome. JMS 2020; 23(3):33-35

Received: 2020-08-26 Accepted: 2020-09-09

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

till 5 years of age when she started with progressive lethargy and slowness of movements. She stopped playing and used to remain in the kitchen for most part of the day. She also developed poor appetite, constipation and puffiness of face for the past 1 year. Her scholastic performance remained average. She was evaluated by a local physician and was advised thyroid function test revealed total T4 of <0.3 mcg/dL and TSH > 150mIU/L. Patient was started on levothyroxine 100mcg tablet. One day after starting levothyroxine she developed vaginal bleeding and was thus referred.

The child was short; she had a puffy face and thick skin. She had no thyromegaly, café-au lait macules or skeletal deformities. She had a pulse rate of 88 beats per minute, regular and normal in volume and character. Her anthropometry revealed height of 106cm (-3.0 SDS; Height age of 5 years), target height SDS was -0.7, and weight of 21 kg (-0.5 SDS). Her arm span was 99cm, US/LS were 53cm/53cm. OFC was 51cm. She had an SMR of B2P1 with no axillary hair. Other systemic examination was unremarkable. There was no delay in tendon reflexes. Her investigations revealed a hemoglobin of 9.4 g%, MCV was 93fL, MCH was 29, TLC was 5,200/mm³, platelet count was 3.17 lakhs, serum total cholesterol was 174mg/dL, triglycerides were 157mg/dL, blood urea 31mg%, serum creatinine 0.56mg%, LFT was normal. The creatinine kinase level was 269, LDH was 377, uric acid was 4.2mg%, serum calcium was 9.47mg%. Hormone profile revealed LH 0.4 IU/L, FSH 3.54 IU/L, Prolactin 54.46 ng/mL, E2 47 pg/ml, 8AM cortisol 8.58 mcg/dL, Total T3 2.20 ng/mL, Total T4 16.7 mcg/dL, TSH 9.31 mIU/L, AntiTPO Ab 370.5 IU/L. X ray knee did not show any epiphyseal changes. Bone age was delayed (Greulich and Pyle method) 3 years (BA << HA < CA). USG pelvis revealed bilateral ovaries showing multiseptate cysts, right ovary volume of 14 cc, left ovary volume of 40cc, endometrial thickness was 4mm, size of the uterus was 5.3x1.8x2.7 cm with an ellipsoid volume of 12.9 mL. USG thyroid revealed that the right lobe of the thyroid was 7*8*16 mm and left lobe was 5*7*10 mm with a heterogeneous echopattern. A final diagnosis of Juvenile autoimmune thyroid disease with primary hypothyroidism and peripheral (GnRH-independent) precocious puberty (Van Wyk -Grumbach Syndrome) was made. The child was asked to continue treatment with 100mcg of levothyroxine and the parents

were reassured. After one year of treatment and follow up, the child is doing well and has not had any further bleeding per vaginum. She has grown in height, her breast regressed to B1 and she also reduced her weight.

DISCUSSION

The eponymous Van Wyk-Grumbach syndrome is one of the most puzzling, yet interesting pediatric complexes. Clinical pointers to differentiate it from true (GnRH dependent) precocious puberty include first; thelarche and menarche without pubarche, second; short stature as opposed to increased height velocity in GnRH dependent PP, and third; delayed bone age compared to height age and both less than chronological age as illustrated in this case.³ LH is prepubertal as is expected in any etiology of GnRH independent PP. It also exemplifies a situation which needs masterly inactivity on the part of the clinician, as the multicystic ovaries found in this condition is important to recognise as a part of this syndrome and thus unnecessary investigation or inadvertent surgical management should be avoided. Also if an MRI (magnetic resonance imaging) of the brain is done as part of evaluation of precocious puberty, it is likely to show a sellar mass which is also to be expected here; but the unwary might seek a neurosurgery consult. This condition incorporates points pertinent to TSH and FSH interaction; it is an example of "hormonal overlap" or "specificity spillover" where high levels of TSH (as in our case) stimulate the FSH receptors in the ovarian granulosa cells to produce estrogen. The role of prolactin in mammary gland and ovary development and physiology has also been postulated as a potential mechanism. Our case illustrates the situation where menstrual bleed happens after the start of treatment for chronic acquired severe hypothyroidism.⁴ It may be prudent to convey this to the parents of such children beforehand, to allay their anxiety later. Also it could be debated whether starting with a lower dose of levothyroxine and slow up-titration might prevent such a sequel in children presenting with a similar clinical picture.⁵

REFERENCES

- [1] Wyk JJV, Grumbach MM. Syndrome of precocious menstruation and galactorrhea in juvenile hypothyroidism: an example of hormonal overlap in pituitary feedback. *J Pediatr* 1960;57:41635.
- [2] Sanjeevaiah AR, Sanjay S, Deepak T, Sharada A, Srikanta SS. Precocious puberty and large multicystic

- ovaries in young girls with primary hypothyroidism. *Endocr Pract* 2007;13:6525.
- [3] Reddy P, Tiwari K, Kulkarni A, Parikh K, Khubchandani R. Van Wyk Grumbach Syndrome: A Rare Consequence of Hypothyroidism. *Indian J Pediatr* 2018;85:102830.
- [4] Zhang H, Geng N, Wang Y, Tian W, Xue F. Van Wyk and Grumbach syndrome: Two case reports and review of the published work. *J Obstet Gynaecol Res* 2014;40:60710.
- [5] Tran S, Kim EE, Chin AC. Severe menorrhagia, unilateral ovarian mass, elevated inhibin levels, and severe hypothyroidism: An unusual presentation of Van Wyk and Grumbach syndrome. *J Pediatr Surg* 2013;48.

SHORT RP TACHYCARDIA, SUCCESSFULLY MANAGED WITH RADIO FREQUENCY ABLATION

Ajaz Ahmad Lone, Nisar Ahmad Tramboo, Syed Bilal

Department of Cardiology, SKIMS

35-year female patient presented with recurrent palpitations. Her baseline ECG was normal but during the episode of palpitations it revealed short RP tachycardia. Intracardiac ECG revealed orthodromic reentrant tachycardia (ORT) due to concealed left lateral pathway (Figure 1). She was successfully treated by radio-frequency ablation (RFA) through trans-septal approach. Figure 2 shows RV pacing getting conducted over left lateral pathway to atria which disappears after ablation of the pathway.

Paroxysmal supraventricular tachycardias (PSVT) are common arrhythmias affecting all ages, patients experience frequent palpitations which may be self-limiting or need management in hospital settings. PSVTs are generally of two types. Those related to and arising within the AV node referred to as AV node reentry tachycardia (AVNRT) and the other type involves a manifest or concealed accessory pathway referred as atrioventricular reentry tachycardia (AVRT). When the antegrade limb of the reentry circuit involves the AV node it presents as a narrow complex tachycardia also called as ORT. Antidromic tachycardia presents as wide QRS tachycardia as the antegrade limb involves the accessory pathway. Typical AVNRT and ORT both present with short RP tachycardia and acute management involves vagal maneuver like Valsalva maneuver and carotid sinus massage. IV medications like adenosine and verapamil or diltiazem frequently terminate this tachycardia. RFA is curative in such arrhythmias with high success rates exceeding 98% and low complication

rates of less than 1%. RFA is indicated in patients presenting with recurrent arrhythmias failing conservative approach or medical treatment.

Fig. 1. Intracardiac ECG showing short RP tachycardia

Fig. 2. RV pacing conducted over left lateral pathway to atria, disappears after ablation of the pathway.

Correspondence:

Dr. Ajaz Ahmad Lone, DM
Additional Professor, Department of Cardiology, SKIMS
Email: drajaz666@gmail.com

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.776>

How to cite this article: Lone A, Tramboo NA, Bilal S. SHORT RP TACHYCARDIA, SUCCESSFULLY MANAGED WITH RADIO FREQUENCY ABLATION. JMS 2020; 23(3):36-37

Received: 2020-08-24 Accepted: 2020-08-29

Lone A et al, Short RP Tachycardia, Successfully Managed with Radio Frequency Ablation.

NASAL RECONSTRUCTION

Peerzada Umar Farooq Baba, Adil Hafeez Wani

Department of Plastic surgery, SKIMS, Soura

SHORT CLINICAL HISTORY AND PROCEDURE:

A 65-year male presented with squamous cell carcinoma of the left side of the nose. Wide local excision of the tumor with reconstruction by ipsilateral forehead flap was done. After 3 weeks flap was detached giving a good functional as well as an aesthetic result.

Written informed consent was taken from the patient.

Squamous Cell Carcinoma on Left Side of Nose

Tumour Excision Performed and Markings for Forehead Flap Done

Nasal Defect Covered by the Flap

Detachment of Flap and Final Insetting

Correspondence:

Dr. Peerzada Umar Farooq Baba, MS
Associate Professor, Department of Plastic surgery, SKIMS
Email: drumar397@gmail.com

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.778>

How to cite this article: Baba PUF, Wani AH. NASAL RECONSTRUCTION. JMS 2020; 23(3):38

Received: 2020-08-28 Accepted: 2020-09-01

LETTER TO EDITOR

Adoption of Telemedicine -need of the hour

Dear Editor,

I would like to describe our experience in telemedicine for the provision of health care in the Northern Territory of Australia and its relevance for the state of Jammu and Kashmir. Northern Territory is a large landmass spread over 1.4 million square kilometers with a population of only 250,000. Major healthcare facilities are located in cities of Darwin and Alice Springs with smaller hospitals in Katherine, Gove, and Tennant Creek. Many small communities are located in remote areas with limited access to health care. We have adopted telemedicine more than a decade ago and gradually scaled up the services which came handy during the latest pandemic due to SARS-CoV-2.

Telemedicine encompasses the delivery of health care using a variety of telecommunication tools. In addition to improved access to health care, telemedicine can reduce the cost and improve convenience. Specialists triage suitable referrals to teleclinics. The patient/carer is informed of the appointment and consent is obtained. The clinic staff procures the relevant investigation reports to be ready at the time of the consult. To run an efficient and effective teleclinic it is essential to have appropriately trained administrative support. Chronic disease management, wherein physical examination is not crucial for decision making, is well suited for telemedicine. Teleclinics are also suitable to assess the need for patients to come into the hospital as well as provide pre-operative advice and preparation prior to planned elective surgical procedures.

Multidisciplinary case conferences with patients, family members, general practitioners, and hospital-based specialists work well to formulate an individualized management plan for complex patients. People living in

remote locations in Jammu and Kashmir are likely to experience difficulty in reaching Srinagar and Jammu due to limited public transportation options, hilly terrain, heavy snow in winter, political upheaval and the frequent imposition of restriction on movement by the law enforcement authorities.

Patient factors that determine the feasibility of telemedicine include access to a telephone (landline or mobile), internet connection with sufficient bandwidth, hearing and communication ability, and ability to use video applications on a mobile phone (smartphone) or computer for video consultations. Many patients in remote communities have low literacy and limited access to internet services in addition to sub-optimal communication skills. These patients access teleclinics by going to the nearest community health center from where they can link up to the health care providers in major centers. We involve interpreter services, family members, health care workers in the community health center in the discussions for patients who have poor health literacy, hearing difficulties, or whose first language is other than English. Seamless link up with pathology, radiology, and pharmacy services to enable electronic test requests, test/imaging reports and prescriptions will improve the client experience.

Telehealth can play a meaningful role in natural disasters and pandemics by helping in “forward triage” before patients arrive in the emergency department as well as directing patients to the hospitals with bed capacity^{1,2}. The provision of telehealth for patients in pandemic lockdown or quarantine ensures delivery of necessary care while protecting health care staff and the community from exposure. We have converted most of the scheduled clinic

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.780>

How to cite this article: Chitturi S. Adoption Of Telemedicine - need of the hour. JMS 2020; 23(3):39-40

Received: 2020-08-31 Accepted: 2020-09-01

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

visits to teleclinic visits allowing for both patients and clinicians to stay at home and provide uninterrupted clinical care. This is important to reduce the emergency department presentations by patients with non-COVID related illnesses.

Innovation and rapid scaling up of telehealth services with improved infrastructure and staff and patient training to support the telehealth service can go a long way to provide more equitable and cost-effective health care to the neediest. Medicare coverage for telehealth services was rapidly expanded in Australia which is likely to have far-reaching effects on reshaping the healthcare landscape. For certain health conditions, telehealth can be the preferred first option with in-person consultations being the second or third option to meet the patient's health care needs³.

REFERENCES

1. Lurie N, Carr BG. The role of telehealth in the medical response to disasters. *JAMA Intern Med* 2018; 178:745-746.
2. Hollander JE, Carr BG. Virtually perfect? Telemedicine for COVID-19. *N Engl J Med* 2020.
3. Duffy S, Lee TH. In-person health care as option B. *N Engl J Med* 2018; 378:104-106.

Dr. Sridhar Chitturi MD, DM, FRACP

Senior Consultant Endocrinologist

Royal Darwin Hospital

Tiwi, NT 0810, Australia

Email: sridharchitturi@gmail.com

Practice Point

High Flow Nasal Cannula Oxygenation

Tajamul Hussain Shah, Suhail Mantoo, Rafi A Jan

Department of Internal and Pulmonary Medicine, SKIMS

The current pandemic of COVID-19 caused by novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) tested the healthcare infrastructure throughout the globe at all possible levels. Early reports suggest about 20% of patients infected with SARS-CoV-2 require hospitalization and 5-8% require admission to intensive care unit (ICU) due to severe disease.^{1,2} Supplementation of oxygen provided by various conventional oxygen therapy (COT) devices (like nasal prongs, face mask, venture mask or non-rebreather mask) may not be sufficient in cases of worsening respiratory failure. One form of escalating respiratory support in such patients is a high flow nasal oxygenation device. High flow nasal cannula oxygenation (HFNC) is a form of non-invasive respiratory support^{3,4}. It acts as a bridge between low flow devices and non-invasive ventilation and may reduce the need for intubation.

Limitations of conventional oxygen supplementation devices^{3,4}

- Due to flow limitation COT devices are not able to provide higher flow required by patients in respiratory failure, resulting in entrainment of room air and reduction in delivered FiO_2 .
- Due to lack of adequate humidification, COT devices provide dry and cool oxygen that can lead to airway inflammation and impaired mucocilliary function which can increase airway resistance and reduced secretion clearance.

High flow nasal cannula (HFNC) oxygenation

HFNC oxygenation device provides high flow oxygenated

gas that is heated and humidified to body condition and is delivered via a special nasal cannula at higher flow rates (10-60 l/min).

Mechanisms of Clinical benefit with HFNO³

- **Heated and humidified air**

In contrast to COT devices, HFNC provides heated and humidified air which helps in maintaining airway hydration, prevents airway desiccation and epithelial injury thereby preserving mucocilliary function.

- **Higher flow rates**

HFNC by virtue of its higher flow rate (~60 L/min) prevents entrainment of room air and therefore provides higher and more reliable FiO_2 . Besides, higher flow rate is also associated with reduction in inspiratory effort, work of breathing and upper airway dead space due to continuous flushing of CO_2 from nasopharynx.

- **Positive end expiratory pressure (PEEP) effect**

HFNC delivers a PEEP of upto 1cm of H_2O for every 10L/min of flow delivered with closed mouth breathing. This PEEP effect can potentially counteract the auto-PEEP, thereby, increasing alveolar recruitment and improving oxygenation.

Advantages of HFNO

HFNC as compared to COT devices and NIV is more comfortable and better tolerated. NIV requires a properly fit

Access this article online

Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v23i3.784>

Correspondence:

Dr. Suhail Mantoo, MD

Assistant Professor, Department of General Medicine, SKIMS

Email: suhail.mantoo@skims.ac.in

How to cite this article: Shah TH, Mantoo S, Jan RA. High Flow Nasal Cannula Oxygenation. JMS 2020; 23(3):41-44

Received: 2020-09-05 Accepted: 2020-09-09

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

interface (mask) which in turn hinders clearance of secretions, oral intake, communication as well as proper positioning during awake self-proning.

Uses of HFNO^{5,6,7}

Oxygenation with HFNC has been shown to have similar and in some cases superior clinical efficacy as compared to COT and NIV. Various clinical scenarios where HFNO has been used include.

- **Management of acute hypoxemic respiratory failure in immunocompromised patients:** Immunocompromised patients with hypoxemic respiratory failure that require intubation have higher mortality. Use of HFNC was associated with significant reduction in rate of intubation and mortality as compared to COT and NIV.
- **Preventing reintubation:** Oxygenation with HFNC reduces the rate of reintubation as compared to COT in low risk medical patients, where as in high risk patients HFNC was non inferior to NIV.
- **Post operative respiratory failure:** HFNC is a reasonable alternative to NIV and COT for treating postoperative respiratory failure and preventing reintubation especially after cardiothoracic, abdominal and lung resection surgeries.
- **Pre-oxygenation:** HFNC has also been used in providing pre-oxygenation to prevent desaturation during endo-tracheal intubation.
- **To facilitate interventions like bronchoscopy:** Patients with hypoxemic respiratory failure are prone to significant desaturation during the bronchoscopy. HFNC can provide respiratory support and reduce the duration and severity of hypoxemia during the procedure.
- **Adult respiratory distress syndrome (ARDS):** Few studies have demonstrated the successful use of HFNC in patients with moderate to severe ARDS.

Precautions

Use of HFNC is considered to be associated with aerosol generation and therefore has potential to spread the infection among health care workers. However use of negative pressure rooms, high energy particulate

accumulator (HEPA) filters, adequate personal protective equipment (PPE) and surgical mask or an N-95 mask over HFN cannula are sufficient to reduce the transmission of infection.⁸

Components of HFNC device and how to set it up^{4,9}

The HFNC apparatus comprises of following components (Fig.1)

- **Humidifier:** Provides heated and humidified gas to the patient through heated inspiratory circuit.
- **Air-Oxygen blender:** At air /oxygen blender, the inspiratory fraction of oxygen (FiO₂) can be adjusted from 21% to 100%.
- **Flow meter:** Allows adjustment of oxygen flow rate upto 60 L/min.
- **Nasal cannula**

Inspiratory circuit: Connects the humidifier with the nasal cannula.

Fig.1: Various components of HFNC device (Source: Wikipedia)

Fig.2: Various adjustable parameters on device (temperature, flow rate and FiO₂).

Setting up of a device

- **Choosing appropriate size cannula:** Nasal cannula should fit well into the nares; however prongs should not totally occlude the nostrils and should allow ample egress of expired gas. A tight fitting nasal cannula can lead to uncomfortably higher distending pressure in nasopharynx.
- **Setting Flow rate:** Maximum physiological benefit of HFNO is achieved at higher flow rates. Although there is no standard way to set the flow rate, an approach that is frequently used in clinical practice is to initiate the flow of 30L/min. Lower flow is preferred in case of COVID-19 to reduce aerosol generation unless higher flows are required to treat respiratory failure.
- **FiO₂:** FiO₂ is set next to achieve a desired peripheral oxygen concentration. **(88-92% in patients underlying COPD, 92-96% for others)**
- **Temperature:** For optimum humidification of air the temperature is set at 37°C. However if there is rain out in the humidifier circuit tubing it may be necessary to reduce the set temperature to 34-35°C.

Patient's respiratory rate (RR), peripheral oxygen saturation (SPO₂) and breathing pattern is monitored continuously. In case there is no improvement in oxygen saturation, reduction in respiratory rate or respiratory distress, flow rate is increased at an increment of 10L/min till desired oxygenation target is achieved. If maximum tolerable flow rate fails to improve oxygenation, FiO₂ is adjusted to achieve same.

Predictors of failure of HFNO

Patients requiring HFNO are at risk of severe respiratory failure, therefore it should be preferably used in areas with facility for close monitoring and intubation if need arises. Indicators of failure include.

- Increasing respiratory rate and heart rate.
- Failure to achieve a target oxygenation despite a high flow rate and FiO₂
- Rising PaCO₂ level.
- Dys-synchronous breathing
- Alteration in mental status and

- Hemodynamic instability.

Use of ROX index (defined as ratio of oxygen saturation as measured by pulse oximetry/FiO₂ to respiratory rate) has been suggested to predict success or failure of HFNC. ROX index more than 4.88 at 12 hours after HFNO predicts success and less likely-hood of intubation where as score less than equal to 3.85 at 12 hour predict failure with 100% specificity.¹⁰

Weaning from HFNO

HFNO is well tolerated and can be used for prolonged periods. Shifting of patient from HFNO to COT can be considered once the flow rate reaches to < 20L/min and FiO₂ of < 50%.

CONCLUSION

HFNO is better tolerated and more comfortable than COT and NIV. Its use may reduce the need of intubation or escalation of respiratory support as compared to COT. Therefore HFNO should be considered in patients with persistent respiratory failure despite COT.

REFERENCES

1. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China. JAMA. 2020;323(13):1239-1242
2. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet 2020; 395: 497506
3. Rali AS, Garies T, Narendra D, Patel P, Guntupalli K. High-flow nasal cannula: COVID 19 and beyond . Ind J Resp Care. 2020;9:134-140
4. Raoof S, Nava S, Carpati C, Hill NS. High-Flow, Noninvasive Ventilation and Awake (Nonintubation) Prone in Patients with COVID-2019 with Respiratory Failure. Chest. 2020; S0012-3692(20)31910-3.
5. Drake MG. High-Flow Nasal Cannula Oxygen in Adults: An Evidence-based Assessment. Ann Am Thorac Soc. 2018; 15(2):145-155.
6. Ding L, Wang L, Ma W, He H. Efficacy and safety of early prone positioning combined with HFNC or NIV in moderate to severe ARDS: a multi-center

- prospective cohort study. Crit Care. 2020; 24(1):28.
7. Messika J, Ben Ahmed K, Gaudry S, et al. Use of High-Flow Nasal Cannula Oxygen Therapy in Subjects with ARDS: A 1-Year Observational Study. Respir Care. 2015; 60(2):162-169.
 8. Li J, Fink JB, Ehrmann S. High-flow nasal cannula for COVID-19 patients: low risk of bio-aerosol dispersion. Eur Respir J. 2020; 55(5):2000892. Published 2020 May 14.
 9. Nishimura M. High-flow nasal cannula oxygen therapy in adults. J Intensive Care. 2015; 3(1):15. Published 2015 Mar 31. doi:10.1186/s40560-015-0084-5
 10. Goh KJ, Chai HZ, Ong TH, Sewa DW, Phua GC, Tan QL. Early prediction of high flow nasal cannula therapy outcomes using a modified ROX index incorporating heart rate. J Intensive Care. 2020; 8:41.

Genetics and the COVID-19 Pandemic

The clinical spectrum of COVID-19-related disease ranges from asymptomatic to severe, fatal presentations.

Older age, gender and comorbidities being identified (hypertension, diabetes, respiratory system disease and cardiovascular disease). Fatal course also presenting in young, previously healthy individuals. Differences in clinical outcomes may be determined by genetic susceptibility.

Sources

INT J INFECT DIS. 2020 MAY;94:91-5.

EUR J CLIN INVEST. 2020 APR 30;10.1111/ECL.13259.

J INFECT. 2020 MAR;80(3):350-371.

COVID-19 and ACE2

Severe lung interstitial edema and leukocyte infiltration in Ace2 KO mice are attenuated in homozygous (Ace^{-/-}) or heterozygous (Ace^{+/-}). Balance of ACE2/ACE levels is the key to lung injury/lung protection during an inflammatory storm. The activation of the RAS axis due to binding of Covid19 to ACE2 partly drives the systemic manifestations of SARS-CoV-2.

Sources

NATURE. 2005 JUL 7;436(7047):112-6.

CIRC RES. 2020 MAY 8;126(10):1456-1474.

TMPRSS2 is a host cell factor critical for spread of several relevant viruses, including influenza A viruses and coronaviruses.

TMPRSS2 is dispensable for development and homeostasis. A TMPRSS2 inhibitor (serine protease inhibitor camostat mesylate) is approved for (unrelated) clinical use, can block virus entry and might constitute a treatment option for current SARS-CoV-2 patients.

Sources

CELL. 2020 APR 16;181(2):271-280.E8.

MOL CELL BIOL. 2006 FEB;26(3):965-75.

ANTIMICROB AGENTS CHEMOTHER. 2016 OCT 21;60(11):6532-6539.

1. The Lancet. [https://doi.org/10.1016/S2213-2600\(20\)30402-1](https://doi.org/10.1016/S2213-2600(20)30402-1)

2. Journal of Burn Care and Research.

<https://doi.org/10.1093/jbcr/iraa034>

3. Journal of the American Medical Association (JAMA). https://www.medicatradefair.com/en/News/News_from_the_Editors/Blood_test_shows_promise_in_the_diagnosis_of_Alzheimer_s

4. Elsevier : Archives of Medical Research. <https://www.sciencedirect.com/science/article/abs/pii/S0188440919310045>

5. The New England Journal of Medicine. <https://www.nejm.org/doi/full/10.1056/NEJMoa2026920>

6. Scientific Reports. <https://www.nature.com/articles/s41598-020-70086-y>

7. Journal of Orthopaedic Surgery and Research. <https://josr-online.biomedcentral.com/articles/10.1186/s13018-020-01930-0>

8. Clinics in Chest Medicine. <https://pubmed.ncbi.nlm.nih.gov/32800203/>

9. Nature Reviews Gastroenterology & Hepatology. <https://www.nature.com/articles/s41575-020-00363-8>

10. The New England Journal of Medicine. <https://www.nejm.org/doi/full/10.1056/NEJMoa2019422>

11. Journal of Paediatric Infectious Diseases. <https://pubmed.ncbi.nlm.nih.gov/29447395/>

Journal of Medical Sciences (JMS) Call for Papers

Articles are invited for publication in Journal of Medical Sciences (JMS); the official journal of Sher-i-Kashmir Institute of Medical Sciences (SKIMS) Srinagar. JMS is a peer-reviewed multidisciplinary medical journal and publishes articles in all branches of clinical medicine and surgery, and allied specialities with particular emphasis on original research articles of regional interest. Currently the Journal is published biannually; the online issue which has been started recently is expected to be published more frequently.

JMS is intended for medical practitioners working in all clinical setting including primary and secondary care. Articles are mainly aimed at non-specialist hospital doctors, family physicians and postgraduate teaching.

JMS invites following types of articles for publication

1. Editorials & Commentaries (by invitation only)
2. Original research paper
3. Review articles (by invitation only)
4. Short article (Brief Report/Mini-Review)
5. Case Series
6. Case Reports
7. Clinical Images
8. Education & Teaching related (Practice point)
9. Letters
10. News & Views

Advantages of publishing in JMS

1. Online Submission: Fast and easy online submission.
2. Fast Peer Review: give important feedback on getting the paper better.
3. International Editorial Board.
4. Support for statistical analysis of the submitted paper.
5. Free online access to all JMS articles
6. Indexed in 'Google'; indexing in other major electronic biomedical databases in process

Please visit our website for Guidance to authors about the required format of the various types of the articles. <http://www.jmsskims.org/instruction-for-authors/>

Thanks
JMS Editorial Team