

Practice Point

Impact of Russian Poplar Allergy in Kashmir: Is It a Call for Action?

Tabasum Shafi, Roohi Rasool

Department of Immunology & Molecular Medicine, SKIMS, Srinagar- 190011

INTRODUCTION

Allergic diseases, including allergic rhinitis (AR) and asthma, are among the fastest-growing chronic disorders, with rising prevalence globally due to urbanisation, pollution, and climate change¹. AR impacts 10–40% worldwide², while asthma is the second leading cause of mortality among chronic respiratory illnesses, affecting 1% and 29% of the population across various countries³. In India, AR prevalence is estimated at 22% 30% and asthma affects over 30 million people⁴. These conditions impair quality of life with elevated healthcare costs, necessitating urgent public health interventions.

The Kashmir Valley has experienced a surge in respiratory allergies in recent decades, with anecdotal reports associating this increase with the widespread release of cottony fluff by the Russian poplar (*Populus deltoides*). These trees, introduced for their economic value in the timber and plywood industries, have raised serious concerns due to their potential to exacerbate seasonal allergies. These trees release pollen during early spring (March–May), coinciding with a surge in OPD visits for allergic rhinitis, conjunctivitis, and asthma exacerbations. The issue also gained public and governmental attention back in 2015, following legal interventions and expert committee recommendations⁵.

Historical and Ecological Context

Russian poplars imported from the United States were introduced into Kashmir in the early 1980s as a fast-growing species capable of meeting the burgeoning timber

demand of the valley. Their short rotation cycle (8–10 years) made them a suitable alternative to Kashmir poplar, which takes 30–40 years to grow. However, the uncontrolled propagation, especially of female clones that produce seed-laden cotton-like fluff, has led to a seasonal environmental and health concern⁶.

Allergic Potential of Poplar Pollen: Evidence and Limitations

Pollen is the principal aeroallergen responsible for initiating allergic symptoms in genetically susceptible individuals. However, not all pollen grains are equally capable of causing allergic reactions in the lower airways. Poplar pollen grains measure ~ 25–40 µm in diameter and are considerably larger than the aerodynamic threshold (<10 µm) required to reach the lower respiratory tract. These grains are generally confined to the upper airways, where they may provoke symptoms such as sneezing, nasal congestion, and eye irritation. However, their role in eliciting lower airway symptoms such as life-threatening asthma is considered scientifically unsupported^{7,8}.

According to the 2015 report by SKIMS, the role of poplar pollen in allergy is real but often overestimated with approximately 20% sensitization to poplar pollen, based on skin prick testing; however, no mono-sensitization to poplar alone was reported, with most patients showing polysensitization to other aeroallergens such as grasses,

Access this article online



Website:
www.jmsskims.org

DOI:
<https://doi.org/10.33883/jms.v28i2.1451>

Correspondence:

Dr. Roohi Rasool, MD

Professor & Head, Department of Immunology & Molecular Medicine, SKIMS

Email: roohi_wani@yahoo.com

How to cite this article: Rasool R, Shafi T. Impact of Russian Poplar Allergy in Kashmir: Is It Really a Call for Action? JMS 2025; 28(2):56-59

Received: 2025-06-03 Accepted: 2025-06-30



This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

dust mites, pines, and fungal spores⁹. These findings highlight the importance of context-specific diagnostics and public education, discouraging the attribution of springtime allergies to Russian poplar alone.

Misconceptions and Public Perception

A significant barrier to the management of seasonal allergies in Kashmir lies in the widespread public misconception regarding the role of Russian poplar. The most visible aspect of these trees during spring, the release of fluffy, white seed hairs (commonly referred to as poplar fluff) from the female poplar, is often mistakenly assumed to be the primary cause of allergic reactions. However, scientific evidence establishes that the fluff is composed of seed hairs, not pollen grains, and is biologically non-allergenic. Its role is more as a passive carrier rather than an active allergen. Despite this, public anxiety is frequently fuelled by the visual impact of dense seed dispersal, which coincides with a seasonal rise in allergic symptoms due to a multitude of other aeroallergens. The fluff may incidentally carry pollen, fungal spores, and particulate matter, but it does not actively trigger allergic reactions on its own^{10,11}.

This confusion has led to emotionally driven demands for large-scale tree removal, without understanding of the environmental consequences or the true aetiology of seasonal allergies. Such reactions risk overshadowing the need for comprehensive aeroallergen mapping, public education, and scientifically sound forestry policies. A public health communication strategy is thus essential to dispel myths and reorient focus toward evidence-based interventions.

The way forward: Diagnostic Approaches and Allergen Profiling

The SKIMS Immunology department employs a standardised Skin prick test (SPT) panel which includes allergens across multiple categories: 22 pollens, 3 fungi, 3 dust mites, 5 dust mixtures, 4 epithelia, and 5 insects. SPT remains the first-line diagnostic tool in suspected IgE-mediated allergic conditions due to its high sensitivity (80–97%) and specificity (70–95%)¹². Additionally, total serum IgE testing is performed to complement SPT findings. Such diagnostics are essential in distinguishing genuine sensitisation from assumed triggers like poplar fluff, which is non-allergenic but may carry other allergenic particles.

Environmental and Urban Planning:

Poplar trees are frequently found near schools, hospitals, and densely populated urban areas. The absence of regulatory guidelines on the plantation density or spatial zoning of allergenic species has led to unchecked growth. The environmental cost of unregulated plantations includes not only public health risks but also ecological imbalances due to the displacement of native tree species.

Climate Change and Allergy Burden

Global warming, rising temperatures, changing precipitation patterns, and environmental degradation have led to increased production and prolonged presence of airborne allergens^{1,13}. This is reflected in early and prolonged pollen seasons, increased severity and frequency of seasonal allergy symptoms and interaction with other environmental stressors like crop burning, wildfire smoke, and particulate pollution. These changes exacerbate allergic diseases and place an additional burden on healthcare systems.

Recommendations and Interventions

Drawing on the findings of the 2015 SKIMS advisory report, expert clinical inputs, and public health practice insights, the following comprehensive and practical recommendations are proposed for managing the health and environmental implications of Russian poplar allergy in Kashmir:

A. Public Health and Education

1. **Public Awareness Campaigns:** A concerted effort should be made to inform the general population that the cotton-like material dispersed by female poplars during the spring season is not an allergen itself. However, it may carry fungal spores and pollen that can exacerbate allergic conditions. Public health departments should initiate community awareness drives during the seed dispersal period (April–May), recommending the use of protective measures such as masks and avoiding peak exposure times.
2. **Clinical Infrastructure and Medical Training:** Strengthen diagnostic services at primary and secondary healthcare levels through the provision of standardised allergy testing (e.g., skin prick testing and serum IgE analysis). Simultaneously, continuing

medical education (CME) modules should be offered to physicians on the clinical recognition, differential diagnosis, and management of aeroallergen-related illnesses.

B. Regulatory and Environmental Management

1. **Phased Removal of Female Poplars:** The government should formulate and enforce legislation for the gradual removal of female poplar trees, especially from sensitive areas such as schools, hospitals, residential zones, tourist destinations, and along riverbanks and flood channels. Trees should be removed or harvested by the 8–10 year growth stage, before entering the reproductive phase, to prevent seed dispersal.
2. **Looping of Female Tree Branches:** In areas where immediate felling is not feasible, looping (pruning) of upper branches during late winter (February–March) should be mandated. This technique has been shown to significantly reduce seed output by limiting the number of flowering catkins.
3. **Urban Forestry Guidelines:** Urban development authorities should integrate allergenic vegetation control into land use planning. Female poplars should not be planted near high-occupancy public spaces. Instead, native non-allergenic or low-allergenic species should be prioritised to ensure ecological balance and public safety.

C. Forestry and Biotechnological Innovation

1. **Promotion of Male Poplar Clones:** Male poplar clones, identifiable during the flowering season by their visible catkins, should be marked, propagated, and mass planted as an alternative to female clones. Male trees do not release seed fluff and are therefore more suitable for urban and peri-urban environments.
2. **Early Sex Determination through Biotechnology:** Invest in the development of biotechnological tools that enable early differentiation of male and female saplings using genetic markers. This will allow for informed selection during plantation planning and nursery propagation.
3. **Establishment of a Populatum:** A designated conservatory (populatum) should be established to house and study native, hybrid, and introduced poplar species. The goal is to support comparative ecological

studies and ultimately breed a fast-growing, climate-resilient, and low-cotton-producing hybrid in collaboration with forestry and plant genetics institutes.

D. Research and Integrated Policy

1. **Scientific Surveillance and Pollen Mapping:** Establish real-time pollen monitoring stations across Kashmir to track allergen trends and issue seasonal forecasts. Data should feed into the development of a region-specific pollen calendar to guide both clinicians and sensitive individuals.
2. **District-wide Allergy Prevalence Surveys:** Conduct multicentric, evidence-based epidemiological studies across all districts of Kashmir. These studies should include skin prick testing and serological assessments to provide accurate data on the prevalence and distribution of pollen allergy and co-sensitisations.
3. **Integrated Policy Framework:** Form a multi-sectoral task force with representation from health, forestry, education, municipal administration, and urban planning departments. This body should oversee the implementation of allergen control measures, plantation regulation, and public health preparedness during the pollen season.

CONCLUSION

While the cotton seeds of Russian poplar are not the primary allergen affecting Kashmiri populations, their role as a contributing factor to seasonal allergic diseases cannot be ignored. The current report supports a balanced, evidence-based response involving public education, scientific monitoring, and responsible forestry management. Blanket removal of these trees, without a replacement strategy, may invite ecological and economic setbacks.

A multi-sectoral approach involving health, environment, education, and urban development sectors is essential. The call for action is not one of eradication, but of rational management guided by science, sustainability, and public health priorities.

REFERENCES

1. Damialis A, Traidl-Hoffmann C, Treudler R. Climate Change and Pollen Allergies. In: Marselle MR, Stadler J, Korn H, Irvine KN, Bonn A, editors. Biodiversity and Health in the Face of Climate Change [Internet]. Cham:

- Springer International Publishing; 2019 [cited 2025 Jun 9]. p. 47–66.
2. Kogias C, Drylli A, Panagiotakos D, Douros K, Antonogeorgos G. Allergic Rhinitis Systematic Review Shows the Trends in Prevalence in Children and Adolescents in Greece since 1990. *Allergies*. 2023 Dec;3(4):220–8.
 3. Yuan L, Tao J, Wang J, She W, Zou Y, Li R, et al. Global, regional, national burden of asthma from 1990 to 2021, with projections of incidence to 2050: a systematic analysis of the Global Burden of Disease Study 2021. *eClinicalMedicine* [Internet]. 2025 Feb 1 [cited 2025 Jun 8];80.
 4. Mahesh PA, Moitra S, Mabalirajan U, Garg M, Malamardi S, Vedanthan PK, et al. Allergic diseases in India – Prevalence, risk factors and current challenges. *Clinical & Experimental Allergy*. 2023;53(3):276–94.
 5. Wani AS. Greater Kashmir. 2024 [cited 2025 Jun 9]. Are Russian Poplars a Menace in Kashmir? Available from: <https://www.greaterkashmir.com/editorial-page-3/are-russian-poplars-a-menace-in-kashmir/>
 6. Chavan S, Keerthika A, Bhat S, Rajarajan K, Handa AK, Ahmad S. Poplar (*Populus deltoides*) in Jammu and Kashmir, India: facts and fiction. *Current science*. 2020 Sep 25;119:910–1.
 7. Samant S, Li JT. Allergen Avoidance: Pollen Avoidance - Tree, Grass, and Weed. In: *Encyclopedia of Medical Immunology* [Internet]. Springer, New York, NY; 2014 [cited 2025 Jun 8]. p. 27–9.
 8. Burge HA. An update on pollen and fungal spore aerobiology. *J Allergy Clin Immunol*. 2002 Oct;110(4):544–52.
 9. migrator. Greater Kashmir. 2019 [cited 2025 Jun 9]. Are poplars responsible for respiratory ailments in Kashmir? Available from: <https://www.greaterkashmir.com/todays-paper/op-ed/are-poplars-responsible-for-respiratory-ailments-in-kashmir/>
 10. Hu Y, Ferguson , David Kay, Bera , Subir, and Li C. Seed hairs of poplar trees as natural airborne pollen trap for allergenic pollen grains. *Grana*. 2008 Sep 1;47(3):241–5.
 11. Ortega-Rosas CI, Meza-Figueroa D, Vidal-Solano JR, González-Grijalva B, Schiavo B. Association of airborne particulate matter with pollen, fungal spores, and allergic symptoms in an arid urbanised area. *Environ Geochem Health*. 2021 May;43(5):1761–82.
 12. Heinzerling L, Mari A, Bergmann KC, Bresciani M, Burbach G, Darsow U, et al. The skin prick test – European standards. *Clinical and Translational Allergy*. 2013;3(1):3.
 13. Singh AB, Mathur C. Climate Change and Pollen Allergy in India and South Asia. *Immunology and Allergy Clinics*. 2021 Feb 1;41(1):33–52.